



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/565, 31/575, 31/57 // (A61K 31/575, 31/57, 31/565) (A61K 31/57, 31/565)	A1	(11) International Publication Number: WO 91/00733 (43) International Publication Date: 24 January 1991 (24.01.91)
(21) International Application Number: PCT/CA90/00212 (22) International Filing Date: 5 July 1990 (05.07.90) (30) Priority data: 376,710 7 July 1989 (07.07.89) US (71) Applicant: ENDORECHERCHE INC. [CA/CA]; 2989 de la Promenade Ste-Foy, Quebec G1W 2J6 (CA). (72) Inventor: LABRIE, Fernand ; 2989 de la Promenade Ste-Foy, Quebec G1W 2J6 (CA). (74) Agent: MITCHELL, Richard, J.; Marks & Clerk, 50 O'Connor Street, Third Floor, P.O. Box 957, Station B, Ottawa, Ontario K1P 5S7 (CA).		(81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BG, BR, CA, CH, CH (European patent), DE*, DE (European patent)*, DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GB, GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, MW, NL, NL (European patent), NO, RO, SD, SE, SE (European patent), SU. Published <i>With international search report.</i> <i>With amended claims.</i>
(54) Title: METHOD OF TREATMENT OF ANDROGEN-RELATED DISEASES (57) Abstract <p>A method of treatment of androgen-related diseases such as prostate cancer in susceptible male animals, including humans, comprises administering novel antiandrogens and/or novel sex steroid biosynthesis inhibitors as part of a combination therapy. Sex steroid biosynthesis inhibitors, especially those capable of inhibiting conversion of dehydroepiandrosterone (DHEA) or 4-androstenedione (Δ^4-dione) to natural sex steroids (and testosterone into dihydrotestosterone) in peripheral tissues, are used in combination with antiandrogens usually after blockade of testicular hormonal secretions. Antiandrogens can also be part of the combination therapy. Pharmaceutical compositions and two, three, four and five component kits are useful for such combination treatment.</p>		

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Monaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	PL	Poland
CA	Canada	JP	Japan	RO	Romania
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
DE	Germany	LU	Luxembourg	TD	Chad
DK	Denmark			TG	Togo
				US	United States of America

METHOD OF TREATMENT OF ANDROGEN-RELATED DISEASES**BACKGROUND OF THE INVENTION**

This invention relates to a method of treatment of androgen-related diseases such as prostate cancer in warm-blooded male animals (including humans) in need of such treatment, and in particular, to a combination therapy comprising administering an antiandrogen in association with an inhibitor of sex steroid biosynthesis to such animals. The invention also includes pharmaceutical compositions and kits useful for such treatment. Androgen-dependent diseases include diseases whose onset, maintenance or progress is, at least in part, dependent upon biological activities induced by androgens (e.g. testosterone and dihydrotestosterone). In one embodiment, the invention provides a treatment of hormone-dependent prostate cancer in warm-blooded male animals which comprises administering both an antiandrogen and at least one inhibitor of sex steroid biosynthesis capable of inhibiting conversion of dehydroepiandrosterone or 4-androstenedione to natural sex steroids in extra-testicular and extra-adrenal tissues.

SUBSTITUTE SHEET

While various investigators have been studying hormone-dependent prostate cancer, none have proposed the combination therapy of this invention.

A.V. Schally et al., Cancer Treatment Reports, 68 (No. 1) 281-289 (1984), summarize the results of animal and clinical studies on growth inhibition of hormone-dependent mammary and prostate tumors by use of analogues of luteinizing hormone-releasing hormones, the so-called LHRH agonists and suggest that LHRH analogs and/or antagonists may have potential for treating breast cancer.

T.W. Redding and A.V. Schally, Proc. Natl Acad. Sci. UA 80, 1459-1462 (1983), relates to inhibition of prostate tumor growth in rats by chronic use of an LHRH agonist, [D-Trp⁶]LHRH.

U.S. Patent Number 4,329,364 relates to use of the antiandrogen, 4'-nitro-3'-trifluoromethyl isobutyranilide for treatment of prostatic cancer.

U.S. Patent Number 4,472,382 relates to treatment of prostate adenocarcinoma, benign prostate hypertrophy and hormone-dependent mammary tumors may with various LHRH agonists and treatment of prostate adenocarcinoma and benign hypertrophy by use of various LHRH agonists and an antiandrogen.

U.S. Patent Number 4,659,695 (Labrie) relates to treatment of prostate cancer in animals whose testicular hormonal secretions are blocked. The method of treatment includes administering an antiandrogen such as flutamide as an inhibitor of sex steroid biosynthesis such as aminoglutethimide and/or ketoconazole.

Some clinical improvement in men with prostate cancer by use of the two LHRH agonists, Buserelin and Leuprolide, is also reported by N. Faure et al. at pages 337-350 and by R.J. Santen et al. at pages 351-364. res-

SUBSTITUTE SHEET

pectively, LHRH and its Analogues - A new Class of Contraceptive and therapeutic Agents (B.H. Vickery and J.J. Nestor, Jr., and E.S.E. Hafez, eds), Lancaster, MTP Press, (1984).

R. Santen et al., The Journal of Steroid Biochemistry, volume 20, no 6B, at page 1375 (1984), relates that the use of ketoconazole in combination with chronic administration of Leuprolide in rodents decreased basal and Leuprolide-stimulated testosterone levels.

One of Applicant's Co-pending U.S. Patent Applications Serial Number 07/321,926 filed March 10, 1989, relates to a combination therapy for treatment of estrogen-related diseases by inhibiting ovarian hormonal secretions and administering an antiestrogen in combination with at least one of several enumerated activity blockers, sex steroid formation inhibitors and the like.

D. Kerle et al., The Journal of Steroid Biochemistry, volume 20, no. 6B, at page 1395 (1984) relates to the combined use of a LHRH analogue and ketoconazole producing objective responses in some prostate cancer patients who have relapsed or failed to respond to treatment with a LHRH analogue alone.

F. Labrie et al., The Prostate, 4, 579-594 (1983), disclose that use of a combination therapy of an LHRH agonist (Buserelin) and an antiandrogen (Anandron) to treat advanced prostate cancer in previously untreated patients effects simultaneous elimination of androgens of both testicular and adrenal origin.

F. Labrie et al., J. Steroid Biochem., 19, 999-1007 (1983), disclose the treatment of prostate cancer by the combined administration of an LHRH agonist and an antiandrogen. Labrie et al. disclose animal and clinical data in support of the proposition that the combined LHRH/antiandrogen treatment neutralizes the stimulatory influence of all androgens on the development and growth of androgen-dependent prostatic cancer.

CONFIDENTIAL SHEET

F. Labrie et al., Abstracts of the 7th International Congress of Endocrinology, Excerpta Medica (1984) at page 98 disclose that treatment of prostate cancer patients with LHRH agonists alone causes a transient increase in serum androgen levels lasting for 5 to 15 days before castration levels are reached. While F. Labrie et al. recommend that orchiectomy, estrogen and LHRH agonists alone should not be further used for treatment of prostate cancer in the absence of a pure antiandrogen, there still is a need for a method of treatment of prostate cancer that effects more complete androgen blockage at the start as well as during the full period of treatment.

There are many data indicating that estrogens have a stimulatory effect on prostatic growth (Lee et al., 1981; J. Androl. 2: 293-299; Belis et al., 1983; J. Androl. 4: 144-149; Walsh and Wilson, 1976; J. Clin. Invest. 57: 1093-1097; De Klerk et al., 1985; Prostate 7, 1-12; Habesucht et al., 1987; Prostate 11: 313-326). Estrogens have also been found to enhance the growth-promoting effect of androgens (Farnsworth, 1969; Invest. Urol. 6: 423-427; Groom et al., 1971; Biochem. J. 122: 125-126; Lee et al., 1973; Steroids 22: 677-683).

Estrogen receptors have been demonstrated in human normal, hyperplastic and cancer prostatic tissue (Mobbs et al., 1989; Proc. 84th Endocrine Soc., Meeting, abst. No. 1410; Mobbs et al., 1983; J. Steroid Biochem. 19, 1279-1290; Wagner et al., 1975; Acta Endocrinol. (Kbh), suppl. 193, 52; and also in laboratory animal prostatic tissue (Swanek et al., 1982; Biochem. Biophys. Res. Commun. 106: 1441-1447).

Moreover, androgen receptor levels were found to be elevated in prostatic tissue of patients treated with estrogen, thus indicating a stimulatory effect of estrogen on the level of androgen receptors in prostatic tissue (Mobbs et al., 1983; J. Ster. Biochem. 19, 1279-1290). A similar stimulatory effect of estrogen has been observed in the dog prostate (Moore et al., 1979; J. Clin. Invest. 63, 351-357).

SUBSTITUTE SHEET

In the prostate as well as in many other tissues, testosterone is irreversibly converted by 5 α -reductase into the more potent androgen dihydrotestosterone (Bruchovsky and Wilson, J. Biol. Chem. 243: 2012-2021, 1968; Wilson, Handbook of Physiology 5 (section 7), pp. 491-508, 1975). Inhibitors of 5 α -reductase have been found to inhibit prostatic growth (Brooks et al., Endocrinology 109: 830, 1981; Brooks et al., Proc. Soc. Exp. Biol. Med. 169: 67, 1982; Brooks et al., Prostate 3: 35, 1982; Wenderoth et al., Endocrinology 113, 569-573, 1983; McConnell et al., J. Urol. 141: 239A, 1989); Stoner, E., Lecture on the role of 5 α -reductase inhibitor in benign prostatic hypertrophy, 84th AUA Annual Meeting, Dallas, May 8th, 1989.

The inhibitory effect of the 5 α -reductase inhibitor Merck L. 652,931 on prostatic and seminal vesicle development in the prepubertal rat was described in Proc. 71st Annual Meeting of Endocrine Society, abst. #1165, p. 314, 1989. The inhibitory effect of MK-906 on dihydrotestosterone formation in men has been described in men by Gormley et al., in Proc. 71st Annual Meeting of Endocrine Society, abst. #1225, p. 329, 1989; Imperato-McGinley et al., in Proc. 71st Annual Meeting of Endocrine Society, abst. #1639, p. 432, 1989; Geller and Franson, in Proc. 71st Annual Meeting of Endocr. Soc., abst. #1640, p. 432, 1989, and Tenover et al., in Proc. 71st Annual Meeting of Endocr. Soc., abst. #583, p. 169, 1989. The activity of the 5 α -reductase inhibitors N,N-diethyl-4-methyl-3-oxo-4-aza-5 α -androstane-17 β -carboxamide (4-MA) and 6-methylene-4-pregnene-3,20-dione (LY207320) has been described by Toomey et al., Proc. 71st Annual Meeting of Endocr. Soc., abst. #1226, p. 329, 1989.

SUBSTITUTE SHEET

SUMMARY OF THE INVENTION

It is an object of the present invention to provide combination therapy for the treatment of prostate cancer wherein the treatment selectively inhibits the formation and/or action of hormones which would otherwise contribute to tumor growth.


It is another object of the invention to provide combination therapy having increased effectiveness in slowing or reversing tumor growth.

It is another object of the invention to provide therapy for treating prostate cancer having significantly reduced frequency of unwanted side effects.

It is a further object of the invention to provide kits having a plurality of active ingredients (with or without diluent or carrier) which, together, may be effectively utilized for carrying out the novel combination therapies of the invention.

It is another object of the invention to provide a novel pharmaceutical composition which is effective, in and of itself, for utilization in a beneficial combination therapy because it includes a plurality of active ingredients which may be utilized in accordance with the invention.

In one aspect, the present invention provides a method for treating prostate cancer in humans or other warm-blooded animals in need of such treatment, said method comprising the steps of blocking androgen receptors by administering a therapeutically effective amount of an antiandrogen having as part of its molecular structure a substituted or unsubstituted androgenic nucleus of the formula:



CHIRCTITITE CLEET

said antiandrogen having as another part of its molecular structure at least one side chain represented by the formula:

$-R^1[-B-R^2-]_x L-G$ wherein said side chain is substituted onto said androgenic nucleus at a position selected from the group consisting of 6α , 7α , 14α , 15α , 16α , 17α and 17β , and wherein:

x is an integer from 0 to 6, wherein at least one of L and G is a polar moiety distanced from said ring carbon by at least three intervening atoms, and wherein:

R^1 and R^2 are independently either absent or selected from the group consisting of straight- or branched-chain alkylene, straight- or branched chain alkynylene, straight- or branched-chain alkenylene, phenylene and fluoro-substituted analogs of the foregoing;

B is either absent or selected from the group consisting of $-O-$, $-S-$, $-Se-$, $-SO-$, $-SO_2-$, $-NR^3-$, $-SiR^3_2-$, $-CR^3OR^3-$, $-NR^3CO-$, $-NR^3CS-$, $-CONR^3-$, $-CSNR^3-$, $-COO-$, $-COS-$, $-SCO-$, $-CSS-$, $-SCS-$, $-OCO-$ and phenylene (R^3 being hydrogen or lower alkyl);

L is either a moiety which together with G , forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of lower alkyl, $-CONR^4-$, $-CSNR^4-$, $-NR^4CO-$, $-NR^4CS-$, $-NR^4CONR^4-$

NR^4

$-NR^4C-NR^4-$, $-SO_2NR^4-$, $-CSS-$, $-SCS-$, $-(NO)R^4-$, $-(PO)R^4-$, $-NR^4COO-$, $-NR^4SO_2-$, $-O-$, $-NR^4-$, $-S-$, $-SO-$ and $-SO_2-$ (R^4 and R^5 being independently selected from the group consisting of hydrogen and lower alkyl; and R^5 being selected from the group consisting of hydrogen, nitrile and nitro); and

G is either a moiety which together with L forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, (C_3-C_6) cyclo-

SUBSTITUTE SHEET

alkyl, bromo(lower)alkyl, chloro(lower)alkyl, fluoro(lower)alkyl, cyano-(lower)alkyl, carboxy(lower)alkyl, (lower)alkoxycarbonyl(lower)alkyl, (C₁-C₁₀)aryl, (C₁-C₁₁)arylalkyl, di(lower)alkylamino(lower)alkyl, fluoro-substituted analogs of the foregoing,

said method of treatment further comprising the step of inhibiting sex steroid formation by administering a therapeutically effective amount of at least one sex steroid formation inhibitor.

In another aspect, the present invention provides a method for treatment of prostate cancer in a human or other warm-blooded animal in need of such treatment, said method comprising the steps of inhibiting sex steroid formation by administering a therapeutically effective amount of an inhibitor of sex steroid formation capable of blocking formation of natural sex steroids from dehydroepiandrosterone and from 4-androstenedione in peripheral tissues (extra-adrenal and extra-testicular), or an inhibitor of sex steroid formation having as part of its molecular structure a substituted or unsubstituted sex-steroid nucleus, and, as another part of its molecular structure at least one side chain of the formula

$-R^1[-B-R^2]_xL-G$ substituted onto a ring atom of said sex steroid nucleus wherein:

x is an integer from 0 to 6, wherein at least one of L and G is a polar moiety distanced from said ring carbon by at least three intervening atoms, and wherein:

R¹ and R² are independently either absent or selected from the group consisting of straight- or branched-chain alkylene, straight- or branched chain alkynylene, straight- or branched-chain alkenylene, phenylene and fluoro-substituted analogs of the foregoing;

B is either absent or selected from the group consisting of -O-, -S-, -Se-, -SO-, -SO₂-, -NR³-, -SiR³₂-, -CR³OR³-, -NR³CO-, -NR³CS-, -CONR³-, -CSNR³-, -COO-, -COS-, -SCO-, -CSS-, -SCS-, -OCO- and phenylene (R³ being hydrogen or lower alkyl);

L is either a moiety which together with G, forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of lower alkyl, -CONR⁴-, -CSNR⁴-, -NR⁴CO-, -NR⁴CS-, -NR⁴CONR⁴-

NR⁵
||

-NR⁵C-NR⁴-, -SO₂NR⁴-, -CSS-, -SCS-, -(NO)R⁴-, -(PO)R⁴-, -NR⁵COO-, -NR⁵SO₂-, -O-, -NR⁴-, -S-, -SO- and -SO₂- (R⁴ and R⁵ being independently selected from the group consisting of hydrogen and lower alkyl; and R⁶ being selected from the group consisting of hydrogen, nitrile and nitro); and

G is either a moiety which together with L forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, (C₂-C₆)cycloalkyl, bromo(lower)alkyl, chloro(lower)alkyl, fluoro(lower)alkyl, cyano(lower)alkyl, carboxy(lower)alkyl, (lower)alkoxycarbonyl (lower)alkyl, (C₆-C₁₀)aryl, (C₇-C₁₁) arylalkyl, di(lower)alkylamino (lower)alkyl, fluoro-substituted analogs of the foregoing;

said method of treatment further comprising administering a therapeutically effective amount of an antiandrogen.

In another aspect, the present invention provides a method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of administering a therapeutically effective amount of an inhibitor of 5 α -reductase activity and administering a therapeutically effective amount of an antiandrogen.

In another aspect, the present invention provides a method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of administering a therapeutically effective amount of an inhibitor of 17β -hydroxysteroid dehydrogenase and administering a therapeutically effective amount of an antiandrogen.

In another aspect, the present invention provides a method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of administering an effective amount of an antiandrogen and at least one sex steroid formation inhibitor from the groups consisting of inhibitors of extragonadal and inhibitors of extraadrenal sex steroid formation.

The invention also provides kits or single packages combining two or more active ingredients useful in treating prostate cancer. A kit may provide (alone or in combination with a pharmaceutically acceptable diluent or carrier), an antiandrogen and at least one additional active ingredient (alone or in combination with diluent or carrier) selected from the group consisting of an LHRH agonist or LHRH antagonist, a sex steroid formation inhibitor (preferably an inhibitor of 5α -reductase activity, an inhibitor of 17β -hydroxysteroid dehydrogenase activity or an inhibitor of 3β -hydroxysteroid dehydrogenase activity) and an antiestrogen.

The foregoing active ingredients may also be mixed in any of the foregoing combinations to form pharmaceutical compositions (with or without diluent or carrier) which, when administered, provide simultaneous administration of a combination of active ingredients resulting in the combination therapy of the invention. Preferably, when LHRH antagonist or agonist is used, it is administered parenterally. For this reason, it may be administered separately in instances where the other active ingredients are formulated for oral ingestion.

The term "sex steroid nucleus" includes estrogenic and androgenic nuclei.

As used herein, the term "androgenic nucleus" includes any compound which, in the absence of the side chain substituent specified herein ($R^1[-B-R^2-]_x L-G$), is capable of acting as an androgen as determined by a weight increase of at least 35 percent over a seven-day period of the ventral prostate of castrated rats treated with the compound in question (15 milligrams twice daily per 100 grams of body weight) versus a control group of castrated rats. Treatment should start on the day of castration. The precise test, other than any parameters set forth in this paragraph, is that reported in Labrie et al., J. Ster. Biochem. 28, 379-384, 1987.

As used herein, the term "estrogenic nucleus" includes any compound which, in the absence of the side chain substituent specified herein ($R^1[-B-R^2-]_x L-G$), is capable of acting as an estrogen as determined by a weight increase of at least 100 percent over a seven-day period of the uterus of ovariectomized rats treated with the compound in question (0.5 mg twice daily per 100 grams of body weight) versus a control group of ovariectomized rats. Treatment should start on the day of castration. The precise test, other than any parameters set forth in this paragraph, is that reported in Simard et al., Mol. Endocrinol. 2: 775-784 (1988).

The following conventions apply to structural formulae set forth herein. Unless specifically designated to the contrary, substituents may have either α or β stereochemistry or, where valence permits, may represent one substituent in α position and another in β position. Presence of optional double bonds are independent of each other. All structures include salts thereof. Atoms of any sex steroid nucleus for which no sub-

SUBSTITUTE SHEET

stituent is shown or described may optionally be substituted or unsubstituted so long as such substitution does not prevent the nucleus from functioning as a "sex steroid nucleus" as defined herein. Those atoms having a defined substituent may optionally be further substituted by other substituents where their valence permits such further substitution. As used herein, the term "lower", when describing a chemical moiety means a moiety having 8 or fewer atoms. For instance, a "lower alkyl" means a C₁ to C₈ alkyl. Any moiety of more than two atoms may be straight- or branched-chain unless otherwise specified.

The term "sex steroid formation inhibitor" includes both androgen and estrogen formation inhibitors and encompasses any compound which inhibits the biosynthesis of active sex steroids or their precursors. One mechanism by which sex steroid formation inhibitors act is by blocking enzymes which catalyze production of natural sex steroids (e.g. dihydrotestosterone), 17 β -estradiol and androst-5-ene-3 β -17 β -diol or precursors to such sex steroids (e.g. androstenedione). Examples of such sex steroid formation inhibitors are compounds capable of blocking the enzymatic activity of, for example, 5 α -reductase, 3 β -hydroxysteroid dehydrogenase, 17 β -hydroxysteroid dehydrogenase or aromatase.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

In a preferred aspect, a combination therapy for prostate cancer includes administering active ingredients effective to inhibit a variety of different mechanisms which may, directly or indirectly, lead to prostatic cancer growth. Desirably, the inhibition of biological activity which leads to prostatic cancer growth proceeds selectively, without

substantially inhibiting other desirable biological activity. Side effects of the treatment are therefore minimized.

Activation of prostatic androgen receptors stimulates growth of prostatic cancer cells. Growth may be inhibited by blocking these receptors with antiandrogens as explained herein. Growth may also be inhibited by reducing the concentration of androgens available to activate the receptors by administering at least one sex steroid synthesis inhibitor. An inhibitor of 5 α -reductase catalyzes conversion of testosterone to dihydrotestosterone (DHT). This is a particularly preferred sex steroid synthesis inhibitor because it selectively reduces DHT levels without reducing testosterone levels. DHT stimulates prostatic cancer growth to a much greater extent than does testosterone. Also absence of DHT forecloses fewer desirable biological functions than does absence of testosterone. For many patients, blocking of testosterone production is also appropriate.

It is believed that estrogens may also increase prostatic cancer growth. Without intending to be bound by theory, estrogens appear to at least be involved in increasing the number of androgen receptors, and may stimulate prostatic cancer growth directly by binding estrogen receptors. Regardless of the mechanism by which estrogens contribute to prostatic cancer growth, it has now been found that a combination therapy which includes inhibition of estrogen activity can enhance effectiveness of treatment without inhibiting desirable biological functions which, in males, are largely independent of estrogen.

There is shown in Figure 1 a schematic representation of the site(s) of action of various drugs, enzymes and hormones. The following abbreviations are used: ER: estrogen receptor; AR: androgen receptor; DHEA: dehydroepiandrosterone; Δ^5 -diol: androst-5-ene-3 β ,17 β -diol; Δ^4 -dione:

SUBSTITUTE SHEET

androstenedione; DHT: dihydrotestosterone; Anti-A: antiandrogen; Anti-E: antiestrogen; ARO: aromatase; 3β -HSD: 3β -hydroxysteroid dehydrogenase, Δ^5 - Δ^4 isomerase; 17β -HSD: 17β -hydroxysteroid dehydrogenase; 1: antiandrogen; 2: inhibitor of 5α -reductase activity; 3: inhibitor of 17β -hydroxysteroid dehydrogenase activity; 4: antiestrogen; 5: inhibitor of aromatase activity; 6: inhibitor of 3β -HSD activity.

Referring to Figure 1, + means increase in androgen receptor levels. As may be seen from Figure 1, stimulation of the androgen receptor is shown to stimulate prostatic cancer growth, and is therefore to be prevented. In addition, stimulation of the estrogen receptor leads to increased levels of androgen receptors and thus may, in addition, exert direct stimulatory effects on prostatic cancer growth. The action of estrogens is therefore to be prevented. Blockers of sex steroid formation from DHEA and Δ^4 -dic in peripheral tissues does not cause inhibition of adrenal glucocorticoid formation. For example, cortisol and aldosterone production is not inhibited and significant complications which could result from their inhibition are avoided. The desired inhibition of sex steroid formation is thus aimed selectively at androgens and estrogens.

A method of inhibiting activation of the androgen receptor is treatment with an effective antiandrogen compound having an affinity for the receptor site such that it binds to the receptor site and prevents androgens from binding and activating the site. It is important to select antiandrogens which tend to be pure antagonists and which have no agonistic activity. Otherwise, the antiandrogen which blocks the receptor site from androgens, may itself activate the site. Preferred antiandrogens are discussed in detail below. Because it is extremely difficult to block all receptor sites, it is desirable to simultaneously decrease the concentration of androgens available to activate androgen receptors in

SUBSTITUTE SHEET


the prostatic cancer tissue. Hence, it is desirable to inhibit secretion of androgens by the testis. This may be accomplished by a variety of known techniques including but not limited to surgical orchiectomy or by administering LHRH agonists or antagonists. For example, LHRH analogues act in a manner effective to stop the production of bioactive luteinizing hormone, the hormone necessary to cause the testis to produce and secrete androgens and other hormones which may be converted to androgens in peripheral tissues. For some patients, it may be unnecessary to inhibit testicular hormonal secretions where sufficiently potent antiandrogens and sex steroid biosynthesis inhibitors are administered.

As may be seen from the scheme of Figure 1, a number of hormones (especially DHEA and Δ^4 -dione) released by the adrenals may be converted by a variety of biological pathways into androgens and estrogens in peripheral tissues. The most potent androgen produced is DHT. It is therefore highly desirable to include an inhibitor of 5α -reductase which prevents the conversion of testosterone into the more potent androgen DHT.

In peripheral tissues, in addition to DHT, the precursors DHEA and Δ^4 -dione can be converted into the estrogens Δ^3 -diol and estradiol. It is desirable to have an inhibitor of 17β -hydroxysteroid dehydrogenase which prevents the formation of testosterone as well as of Δ^3 -diol and estradiol. In addition, since Δ^4 -androstenedione can be converted into estrone and then to estradiol, it may be useful to block the activity of aromatase, the enzyme responsible for such conversion. Other sex steroid formation inhibitors, such as inhibitors of 3β -HSD can also be used. However, as mentioned earlier, when 3β -HSD is blocked in peripheral tissues, it is also likely that a similar inhibition will take place in the adrenals, thus leading to low secretion of glucocorticoids and mineralocorticoids. When such compounds are used, essential glucocorticoids and sometimes mineralocorticoids should be added back as part of the therapy.


Estrogens, at physiological concentrations, are known to stimulate the growth of the human prostatic cancer cell line LNCaP. This effect of estrogen may be inhibited, however, by antiestrogenic compounds described herein.

In one embodiment, the present invention provides a method of treating prostate cancer comprising the step of administering a therapeutically effective amount of an antiandrogen, and of administering a therapeutically effective amount of an inhibitor of sex steroid formation which has, as part of its molecular structure, a substituted or unsubstituted estrogen nucleus of general structure I:



wherein the dotted lines represent optional pi bonds; and wherein said compound includes as another part of its molecular structure a side chain substitution onto a ring carbon of said general structure I in at least one position selected from the group consisting of 7, 14, 15, 16, 17 (preferably 7 α , 15 α , or 17 α), said side chain being of the formula $-R^1[-B-R^2-]L-G$, as defined above, wherein general structure I further includes at least one substitution selected from the group consisting of 15-halo, 16-halo, a 15,16 bridge atom (preferably carbon), a 14,15 bridge atom (preferably oxygen), and a 16-pi-bonded lower alkyl.

In certain embodiments, the antiandrogen utilized in the present invention may be represented by the general formula:




wherein the dotted lines represent optional double bonds; wherein R^{10} is hydrogen or lower alkyl, R^{13} is absent, hydrogen or methyl in β position,

$R^{17(\alpha)}$ is selected from the group consisting of hydrogen, hydroxyl, lower alkanoyloxy, lower alkyl, lower alkenyl, lower alkynyl, halo(lower)alkyl, halo(lower)alkenyl, halo(lower)alkynyl and fluoro-


(β)

substituted aromatic ring, and a moiety which, together with R^{17} forms



III


$R^{17(\beta)}$ is selected from the group consisting of hydroxyl, (C_1-C_{10}) alkanoyloxy, (C_2-C_7) alkenoyloxy, (C_2-C_7) alkynoyloxy, aroyloxy, alkenoyloxy, cycloalkanyloxy, 1-alkyloxy-alkyloxy, 1-alkyloxycycloalkyloxy, alkylsilyloxy, carboxyl, alkanoyl and a moiety which together with $R^{17(\alpha)}$ forms



III

Antiandrogens useful in the combination therapy of the invention also include but are not limited to flutamide (available from Schering-Plough Corp., Kenilworth, New Jersey, under trade name EULEXIN), Nilutamide (available from Roussel of Paris, France, under trade name ANANDRON), cyproterone acetate (available from Schering AG, Berlin under trade name ANDROCUR), Casodex available from ICI Pharmaceuticals, Macclesfield, England. Preferably, the antiandrogen has, as part of its molecular structure, a substituted or unsubstituted androgenic nucleus, and as another part of its molecular structure, the side-chain $-R'[-B-R^2-]_x L-G$ as defined above. Numerous syntheses of the preferred compounds set forth in the U.S. patent application of Labrie and Mérand entitled "Androgen Derivatives for use in the inhibition of sex steroid activity" which is being executed on even date herewith, the entire disclosure of which is hereby incorporated by reference as though fully set forth herein. A preferred antiandrogen is

EM 101



(x = 10)

which may be synthesized as set forth below.

Example 1

Synthesis of N-butyl, N-methyl-11-(17' β -hydroxy-4'-androsten-3'-on-7' α -yl) undecanamide (EM 101) (5, $x=10$) (Scheme 1)

17 β -acetoxy-7 α -(11'-hydroxy undecanyl)-4-androsten-3-one (2)

Under argon atmosphere, in a flame dried apparatus with magnetic stirrer, a solution of 11-bromo undecanol tetrahydropyranyl ether (25 g, 74 mmol) in anhydrous THF (150 ml) was added dropwise to iodine-activated magnesium (1.9 g). The mixture was kept at room temperature overnight and then was cooled to -30°C and anhydrous cuprous chloride (0.3 g) was added quickly. After 45 min of stirring at this temperature, commercial 4,6-androstadien-17 β -ol-3-one acetate (1) (10 g, 30.5 mmol) in anhydrous THF (100 ml) was added dropwise during 4 h. After 35 min, acetic acid (6 ml) and water (100 ml) was added. The mixture was allowed to reach room temperature and was stirred overnight. Afterwards, the organic compound was extracted with ether (3X). The organic layers were washed with water, dried on magnesium sulfate and evaporated. The residue was dissolved in acetic acid (35 ml) and water (100 ml) and kept 48 h at room temperature. And then, the organic compounds were extracted with ether (3X). The organic layers were washed with saturated sodium bicarbonate solution and water, dried on magnesium sulfate and evaporated. The product was purified by Silica gel dry column chromatography (Kieselgel, 60F254, Merk, 0.063- 0.200 mm, 150 g). Elution with a mixture of methylene chloride and ethyl acetate (20:1 v/v) gave 17 β -acetoxy-7 α -(11'-hydroxy-undecanyl)-4-androsten-3-one (2a, 1.46 g, 2.8 mmol, 9.2%) as a colorless oil: IR ν_{\max} neat 3450, 1740, 1685, 1620 and 1245 cm^{-1} ; NMR 0.84 (s, 3H, 18'-CH₃), 1.21 (s, 3H, 19'-CH₃), 2.05 (s, 3H, OCOCH₃), 3.61 (t, 2H,

SUBSTITUTE SHEET

J=6.59 Hz, H-C.1'), 4.61 (t, 1H, J=7.69 Hz, H-C.17) and 5.73 (s, 1H, H-C.4) and 17 β -acetoxy-7 β -(11'-hydroxy undecanyl)-4-androsten-3-one (2b, 0.9 g, 1.7 mmol, 5.6%) as a colorless oil.

11-(17' β -acetoxy-4'-androsten-3'-on-7' α -yl) undecanoic acid (3)

To 17 β -acetoxy-7 α -(11'-hydroxy undecanyl)-4-androsten-3-one (2a, 800 mg, 1.6 mmol) dissolved in acetone (50 ml) and cooled to 0°C was added under stirring during 5 min, a solution of Jones' reagent (8N chromic acid solution) (0.283 ml). After 15 min, isopropanol (0.5 ml) was added followed by water and the mixture was extracted with ethyl acetate (3X). The organic layers were washed with brine, dried on magnesium sulfate and evaporated to dryness under reduced pressure. The crude 11-(17' β -acetoxy-4'-androsten-3'-on-7' α -yl) undecanoic acid (3) (740 mg) was used in the next step without purification.

N-butyl, N-methyl-11-(17' β -acetoxy-4'-androsten-3'-on-7' α -yl) undecanamide (4)

To a solution of the above undecanoic acid derivative 3 (390 mg, 0.78 mmol) in anhydrous methylene chloride (8 ml) cooled at -10°C was added, under stirring, triisobutylamine (240 μ l) and isobutylchloroformate (140 μ l). After 30 min, N-methylbutylamine (1.8 ml) was added and the mixture was stirred at room temperature for 1 h. Methylene chloride was added. The organic solution was washed with 1N hydrochloric acid, water, saturated sodium bicarbonate solution and finally with water, dried on magnesium sulfate and evaporated to dryness. The residue was


chromatographed on silica gel (Kieselgel, 60F254, Merck, 0.063-0.200 mm, 20 g). Elution with a mixture of diethyl ether and methylene chloride (1:20, v/v) gave N-butyl, N-methyl-11-(17' β -acetoxy-4'-androsten-3'-on-7' α -yl) undecanamide 4 (230 mg, 0.39 mmol, 46% for the alcohol (2a)) as a colorless oil; IR ν_{\max} neat 1740, 1680, 1640 and 1240 cm^{-1} ; NMR 0.84 (s, 3H, 18'-CH₃), 0.95 (t, 3H, J=6.93 Hz, N-(CH₂)₃CH₃), 1.21 (s, 3H, 19'-CH₃), 2.04 (s, 3H, OCOCH₃), 2.91 and 2.97 (2s, 3H, N-CH₃), 3.26 and 3.36 (2t, 2H, J=7.86 Hz, N-CH₂C₂H₅), 4.61 (t, 1H, J=8.42 Hz, H-C.17') and 5.72 (s, 1H, H-C.4').

N-butyl, N-methyl-11-(17' β -hydroxy-4'-androsten-3'-on-7' α -yl) undecanamide (5) (EM 101)

The above acetoxy amide 4 (170 mg, 0.29 mmol) was dissolved in methanol (20 ml) and 6% potassium carbonate (2 ml) and heated at 65°C for 200 min. After cooling, acetic acid (1 ml) and water (150 ml) were added and the mixture was extracted with ethyl acetate (3X). The organic layers were washed with water, dried on magnesium sulfate and evaporated to dryness. The residue was purified by Silica gel dry column chromatography (Kieselgel, 60F254, Merk, 0.063-0.200 mm, 20 g). Elution with a mixture of diethyl ether and methylene chloride (1:9, v/v) gave N-butyl-N-methyl-11-(17' β -hydroxy-4'-androsten-3'-on-7' α -yl) undecanamide (EM 101, 94 mg, 0.17 mmol, 58%) as a colorless oil; IR ν_{\max} (neat) 3400, 1670 and 1640 cm^{-1} ; NMR 0.80 (s, 3H, 18'-CH₃), 0.95 (t, 3H, J=6.75 Hz, N-(CH₂)₃CH₃), 1.21 (s, 3H, 19'-CH₃), 2.91 and 2.97 (2s, 3H, N-CH₃), 3.25 and 3.35 (2t, 2H, J=7.3Hz, N-CH₂C₂H₅), 3.67 (t, 1H, J=8.18, H-C.17') and 5.72 (s, 1H, H-C.4').

SUBSTITUTE SHEET

SCHEME I



Sex steroid formation inhibitors useful in the combination therapy of the invention include but are not limited to inhibitors of 5 α -reductase activity, inhibitors of 17 β -hydroxysteroid dehydrogenase activity, inhibitors of 3 β -hydroxysteroid dehydrogenase activity and inhibitors of aromatase activity.

A typically suitable 5 α -reductase inhibitor is MK-906, a product of Merck, Sharp & Dohme (Mc Connell et al., J. Urol. 141: 239A, 1989). Another inhibitor of 5 α -reductase is 17 β -N,N-diethylcarbamoyl-4-methyl-4-aza-5 α -androstan-3-one (4-MA) (Brooks et al., Endocrinology 109: 830, 1981; Liang et al., Endocrinology 112: 1460, 1983). Other 4-azasteroids acting as 5 α -reductase inhibitors can be formed in Liang et al., J. Biol. chem. 259: 734-739, 1984; and in Brooks et al., Steroids 47: 1-19, 1986, 6-methylene-4-pregnene-3,20-dione has also been described as 5 α -reductase inhibitor (Petrov et al., J. Endocrinol. 95: 311-313, 1982). Similar properties have been described for 4-methyl-3-oxo-4-aza-5 α -pregnane-30(s) carboxylate (Kadohama et al., J. Natl. Cancer Inst. 74: 475-486, 1985).


Trilostane and epostane have been described as inhibitors of 3 β -hydroxysteroid dehydrogenase activity (Ernshaw et al., Clin. Endocrinol. 21, 13-21, 1984; Robinson et al., J. Steroid Biochem. 21, 601-605, 1984; Lambert et al., Ann. Clin. Biochem. 23, 225-229, 1986; Potts et al., Steroids 32, 257-267, 1978) and have been successfully used for the treatment of breast cancer in combination with corticosteroids (Beardwell et al., Cancer Chemother. Pharmacol. 10: 158-160, 1983; Williams et al., Cancer Treat. Rep. 71, 1197-1201, 1987).

4-MA, (17 β -N,N-diethylcarbamoyl-4-methyl-4-aza-5 α -androstan-3-one) has been found to inhibit 3 β -hydroxysteroid dehydrogenase activity in granulosa cells (Chan et al., Biochem. Biophys. Res. Commun. 144, 166-171, 1987). Epostane has been shown to inhibit 3 β -hydroxysteroid dehydrogenase activity in pregnant goats (Taylor, J. Endocrinol. 113, 489-493, 1987).


SUBSTITUTE SHEET

Preferred inhibitors of 17 β -hydroxysteroid dehydrogenase activity include but are not limited to:


N-butyl, *N*-methyl-11-(16' α -chloro-3',17' β -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 139").



N-n-butyl-*N*-methyl-11-(16' α -chloro-3',17' α -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 170")



N-n-butyl-*N*-methyl-11-(16' α -bromo-3',17' α -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 171")



Examples of certain synthesis schemes for EM 139, EM 170 and EM 171 are set forth below (see example 2 and schemes 2 and 3). Those of skill in the art will recognize analogous schemes for synthesizing analogous compounds.

Example 2

SYNTHESIS OF PREFERRED SEX STEROID ACTIVITY INHIBITORS

Synthesis of a starting compound, N-n-butyl, N-methyl-11-(3'-benzoyloxy-17'-~~oxo~~-estra-1',3',5'(10')-trien-7' α -yl)undecanamide (14a) (SCHEME 2)

19-nor-testosterone acetate 3-enolacetate (7)

In an apparatus supplied with a drierite drying tube, a solution of 19-nor-testosterone (6) (100 g; 0.365 mole) in acetic anhydride (200 ml), pyridine (32 ml) and acetylchloride (320 ml) was heated at reflux under magnetic stirring, for 3 h and then concentrated to dryness under vacuum. The dry residue was triturated in absolute ethanol, filtered and washed with little portions of absolute ethanol. After drying, 19-nor-testosterone acetate 3-enolacetate was obtained as a white powder (121.4 g, yield 93%) mp. 176-177°C. The structure was confirmed by spectroscopic means.

17 β -acetoxy-estra-4,6-dien-3-one (8)

To a cooled suspension of enolacetate (121 g; 0.337 mole) in a mixture of DMF (330 ml) and water (7.2 ml) at 0°C was added, under nitrogen, over a period of 1 h, N-bromosuccinimide (63 g). The resulting solution was stirred for an additional 0.5 h at 0°C. Then lithium carbonate (60.8 g) and lithium bromide (30.4 g) were added. The mixture was heated at 95°C for 3 h and then poured into 1.7 l of ice-cold water containing 165 ml of glacial acetic acid. After stirring during 15 hours, the crude 17 β -acetoxy-estra-4,6-dien-3-one (8) was filtered, washed with water,

dried in a desiccating apparatus and recrystallized twice from isopropyl ether (72 g, yield 68%, mp 110°C). The structure was confirmed by spectroscopic means.

7 α -(11'-acetoxy-undecyl) 17 β -acetoxy estro-4-en-3-one (9)

A. Preparation of reagents and solvents

11-bromo undecanol tetrahydro pyranyl ether

11-bromo-undecanol (100 g, 398 mmol) was dissolved in dry ether (768 ml) and the solution was cooled to 0°C using an ice/H₂O bath. To this solution was added HCl gas (2.13 g, 58.4 mmol, 26 ml of HCl/ether).

To this mixture, a solution of 3,4-dihydro-2H-pyran (39.9 g, 43.3 ml) freshly distilled in dry ether (218 ml) was added over a period of 90 min. The solution was then stirred over a period of 16 hours at room temperature. Afterwards, sodium bicarbonate was added to the mixture. The residue was filtered and the solvent was evaporated under vacuum.

The product was then filtered through basic alumina (250 g, Woelm, grade II) using petroleum ether (30-60) as solvent (112 g, 81%).

B. Grignard reagent

In a dry three-neck flask (1000 ml) under dry argon, magnesium (12.0 g, 494 mmol) was placed and activated with iodine. Magnesium was heated with the flame to remove iodine and to dry the apparatus. The system was then cooled to -20°C, and a solution of 11-bromo-undecanol tetrahydro pyranyl ether (73.8 g, 211 mmol) in dry THF (420 ml) was added dropwise. The mixture was stirred under dry argon during one day at -20°C.

The mixture was cooled to -35°C ($\pm 2^\circ\text{C}$) using a dry ice/CCL₄/acetone bath. The anhydrous cuprous chloride (1.18 g, 12 mmol) was added and the mixture was stirred over a period of 0.5 h.

C. Addition of Grignard reagent

After 0.5 h, using the same apparatus mentioned above (Ar, -35°C), a solution of 17 β -acetoxy estra-4,6-diene-3-one (8) (32.0 g, 102 mmol) in dry THF (300 ml) was added dropwise over a period of 6 h to the Grignard reagent (red coloration appeared and disappeared). The mixture was stirred for an additional 1 h and, after removal the cooling bath, acidified (about 0°C) with acetic acid (40 ml), diluted with water and extracted with ether (3x). The ether solution was washed with a saturated sodium bicarbonate solution and water. The organic layer was dried over anhydrous magnesium sulfate and evaporated under reduced pressure to dryness.

The residue was dissolved in MeOH (660 ml) and 5N HCl (180 ml), refluxed for 1 h and 45 min, then concentrated under reduced pressure and cooled in an ice bath. The mixture was then filtered to remove the white precipitate. After the solution had been diluted with water and extracted with methylene chloride (3x), the organic layer was dried over anhydrous MgSO_4 and evaporated under reduced pressure to dryness. Finally, the product (55.9 g, brown oil) was chromatographed on silica gel (Kieselgel 60F254, Merck, 0.063-0.200 mm, 1500 g). Elution with mixtures of methylene chloride and ethyl acetate (4:1 to 1:2 v/v) and then pure ethyl acetate gave crude 7 α -(11'-hydroxy-undecyl)-17 β -hydroxy estra-4-en-3-one (34.8 g) which was dissolved in dry pyridine (200 ml) and dry acetic anhydride (200 ml), stirred 17 h at room temperature and then poured in ice-water. The product was extracted with methylene chloride (3X), washed with 1N hydrochloric acid, water, saturated sodium bicarbonate and water (3X), dried on anhydrous magnesium sulfate and filtered. After evaporation of solvent, the mixture (35 g) of 7 α - and 7 β -diacetoxyenones and degradation products of Grignard reagent were separated by flash chromatography on

SUBSTITUTE SHEET

silica gel (Kieselgel 60, Merck, 230 mesh ASTM, 2.0 kg) developed with a mixture of hexane and diethyl ether (2:3 v/v). The first product eluted was pure amorphous 7 α -(11'-acetoxy undecyl) 17 β -acetoxy-estra-4-en-3-one, (9) (20.8 g, 39.4 mmol, yield from dienone was 39.0%). Further elution gave the 7 β -isomer (10) (5.4 g, 10.3 mmol, 10%). All structures were determined by spectroscopic means.

7 α -(11'-hydroxy-undecyl) estra-1,3,5(10)-trien-3,17 β -diol (11a)

Under dry argon, a solution of 7 α -(11'-acetoxy undecyl) 17 β -acetoxy-estra-4-en-3-one (9) (17.0 g, 32.4 mmol) in dry acetonitrile (150 ml) was added rapidly to a suspension of cupric bromide (14.8 g, 66.2 mmol) and mmol) and lithium bromide (2.89 g, 33.6 mmol) in warm acetonitrile (75 ml). The mixture was heated to reflux over a period of 30 min and stirred vigorously, and then cooled to room temperature. A saturated aqueous solution of sodium bicarbonate (50 ml) was added, and then the organic compound was extracted with ethyl acetate (3x 150 ml). The organic layers were washed with water, dried over anhydrous magnesium sulfate, filtered and evaporated under vacuum to dryness. The residue was chromatographed on silica gel (Kieselgel 60F254 Merck 0.063-0.200 mm; 1000 g). Elution with hexane-ethyl acetate (1:1 v/v) gave the 7 α -(11'-acetoxy-undecyl) estra-1',3',5'(10')-trien-3,17 β -diol, 17 β -acetate (11b) (8.51 g; 50.3%) and the starting product (1.33 g; 15%).

The above diacetate phenol (8.51 g, 16.2 mmol) was dissolved in methanol (90 ml) and sodium hydroxyde 30% (w/v) (9 ml). The mixture was refluxed for 90 min under dry nitrogen. The solution was then concentrated under vacuum and diluted with hydrochloric acid (10% v/v). The

mixture was extracted using ethyl acetate (4 x 150 ml) and the ethyl acetate extract was washed with water, dried over anhydrous magnesium sulfate, filtered and evaporated under vacuum. The evaporation gave 7 α -(11'-hydroxy undecyl) estra-1,3,5(10)-trien-3,17 β -diol (11a) (6.99 g, 98% brut) as a yellow foam, the structure of which was confirmed by spectroscopic means.

3-benzoyloxy 7 α -(11'-hydroxy undecyl) estra-1,3,5(10)-trien-17 β -ol
(12)

The above triol (6.99 g; 15.8 mmol) was dissolved in acetone (25 ml) and an aqueous solution of sodium hydroxyde (1N, 19.1 ml). The mixture was cooled to 0°C using an ice/water bath. Benzoyl chloride (2.22 ml, 19.1 mmol) was then added dropwise. The mixture was stirred for 40 min at 0°C and then diluted with water. The solution was extracted using ethyl acetate (3X) and the organic layers were washed with a saturated aqueous solution of sodium bicarbonate and finally with water. The ethyl acetate solution was dried over anhydrous magnesium sulfate, filtered and evaporated under vacuum to dryness. Then, the residue was immediately chromatographed on silica gel (Kieselgel, 60F254, 0.063-0.200 mm; 500 g). The chromatography was carried out, first, using methylene chloride as solvent (about 1 liter) and secondly the pure 3-benzoyloxy 7 α -(11'-hydroxy undecyl) estra-1,3,5(10)-trien-17 β -ol (12), colorless oil (6.50 g, 75%) was eluted with methylene chloride-ethyl acetate (5:1 about 1 liter and 4:1; v/v). The structure was confirmed by spectroscopic means.


SUBSTITUTE SHEET

11-(3'-benzoyloxy-17'-oxo-estra-1',3',5'(10')-trien-7' α -yl) undecanoic acid (13)

To a cooled solution of 3-benzoyloxy-7 α -(11'-hydroxy undecyl)estra-1,3,5(10)-trien-17 β -ol (12) (4.3 g) in acetone (100 ml) was added dropwise Jones' reagent (8N-chromic acid solution, 6.7 ml). After 30 min, isopropanol (40 ml) was added and the mixture was concentrated under vacuo. Water was added and the mixture was extracted four times with ethyl acetate. The organic layers were washed twice with brine, dried over magnesium sulfate and evaporated to dryness. The crude 11-(3'-benzoyloxy-17'-oxo-estra-1',3',5'(10')-trien-7' α -yl) undecanoic acid (13) (3.94 g) was used in the next step without purification.

31

Scheme 2



SUBSTITUTE SHEET

N-n-butyl, n-methyl-11-(3'-hydroxy-17'-oxo-estra-1',3',5'(10')-trien-7'- α -yl) undecanamide (14b)

To 11-(3'-benzoyloxy-17'-oxo-estra-1',3',5'(10')-trien-7'- α -yl) undecanoic acid (13) (3.94 g, 7.22 mmol), dissolved in anhydrous CH_2Cl_2 (100 ml) and cooled at -10°C was added tributylamine (2.18 ml, 9.15 mmol) and isobutylchloroformate (1.30 ml, 10.0 mmol). The solution was stirred during 35 min. and N-methylbutylamine (13 ml, 109.7 mmol) was added. The mixture was warmed to room temperature and stirred during 1 h. Afterward, CH_2Cl_2 was added and the organic phase was washed with 1N HCl, water, saturated sodium bicarbonate solution and finally with water, dried with anhydrous MgSO_4 and the solvent was removed under reduced pressure. The residue was purified by chromatography on silica gel. Elution with mixture of EtOAc/hexane (1.5:8.5 v/v) yielded N-butyl, N-methyl-11-(3'-benzoyloxy-17'-oxo-estra-1',3',5'(10')-trien-7'- α -yl) undecanamide (14a) (4.25 g, 96%) as colorless oil; IR ν (neat) 1750, 1725 and 1640 cm^{-1} . The above described benzoyloxy amide (341 mg, 0.54 mmol) was dissolved in methanol (10 ml) and cooled at 0°C . Following this 2N NaOH (5 ml) was added and the mixture was stirred during 60 min. at 0°C . The solution was neutralized with 1N HCl and extracted with CH_2Cl_2 . The organic phase was dried with anhydrous MgSO_4 and the solvent was removed under reduced pressure. The residue was purified by chromatography on silica gel. Elution with mixture of EtOAc/hexane (3:7 v/v) yielded N-butyl, N-methyl-11-(3'-hydroxy-17'-oxo-estra-1',3',4'(10)-trien-7'- α -yl) undecanamide (14b) (284 mg, 97%) as colorless oil; $^1\text{H-NMR}$ δ (CDCl_3) 0.91 (s, 3H, 18'- CH_3), 2.76 app(d, 1H, $J=16.3\text{Hz}$, part of ABX system, 6'-H) 2.96 and 2.98 (2s, 3H N- CH_3), 3.27 and 3.38 (2t_{app}, 2H, $J=7.5\text{Hz}$, N- CH_2 -), 6.63 (broad s, 1H, 4'-H), 6.70 (broad d, 1H, $J=8.5\text{Hz}$, 2'-H), 7.12 (d, 1H, $J=8.4\text{Hz}$, 1'-H); IR ν_{max} (neat) 3270, 1730, 1615 cm^{-1} ; MS m/e 523 (M^+ , 100%), 508 (M^+-CH_3 , 32%), 142 ($\text{C}_7\text{H}_5\text{CON}(\text{CH}_3)\text{C}_4\text{H}_9$, 47%).

SUBSTITUTE SHEET

16-HALO-ESTRADIOL UNDECANAMIDE (SCHEME 3)

N-*n*-butyl, *N*-methyl-11-(3',17'-diacetoxy-estra-1',3',5'(10'),16'-tetraen-7'α-yl) undecanamide (15)

The ketone amide 14b (163 mg, 0.50 mmol) was dissolved in isoprenyl acetate (10 ml). *p*-toluenesulfonic acid (44 mg) was then added and the solution was distilled to about two-thirds of the original volume in 7 h and was then stirred at reflux for 12 h. Afterwards, the solution was cooled with an ice-water bath and extracted with 50 ml of cooled ether. The ether was washed with a cooled saturated sodium bicarbonate and water. The organic phase was dried with anhydrous MgSO₄ and the solvent was removed under reduced pressure. The residue was filtered through alumina (15mm x 50 mm alumina Woehlm neutral, activity II) using a mixture of benzene-diethyl ether (3:7 v/v) as eluant. The solvent was removed under reduced pressure and, the residue was purified by flash chromatography on silica gel. Elution with mixture of EtOAc/hexane (1:4 v/v) yielded the *N*-butyl, *N*-methyl-11-(3',17'-diacetoxy-estra-1',3',5'(10'),16'-tetraen-7'α-yl) undecanamide (15) (244 mg, 80%) as colorless oil; ¹H-NMR δ_m (CDCl₃) 0.92 (s,3H,18'-CH₃), 0.92 and 0.95 (2t,3H,J=7.0 Hz,N(CH₃),CH₃), 2.18 (s,3H,17'-OCOCH₃), 2.28(s,3H,3'-OCOCH₃), 2.76 app (d,1H,J=16.1 Hz, part of ABX system,6'-H), 2.90 and 2.96 (2s,3H,N-CH₃), 3.26 and 3.35 (2t_{app},2H,J=7.6 Hz,N-CH₃-), 5.52 (m,1H,16'-H), 6.80 (broad s,1H,4'-H), 6.85 (dd,1H,J₁=9.1 Hz and J₂=3.0 Hz,2'-H), 7.27 (d,1H,J=9.1 Hz,1'-H); IR ν_{max} (neat) 1750, 1635, 1200 cm⁻¹; MS m/e 607 (M⁺,2%), 5(M⁺-COCH₃,100%),

SUBSTITUTE SHEET

550 ($M^+ - COCH_3 - CH_3$, 13%), 523 ($M^+ - 2COCH_3$, 45%), 142 ($C_8H_8CON(CH_3)C_8H_8$, 55%), 129 ($C_8H_8(CH_3)NCOCH_3$, 38%), 114 ($C_8H_8(CH_3)NCO$, 60%), 86 ($C_8H_8(CH_3)N$, 25%); EXACT MASS calcd for $C_{28}H_{38}O_2N$ 607.4239, found 607.4234.

N-butyl, *N*-methyl-11-(16' α -chloro-3'-acetoxy-17'-oxo-estra-1',3',4'-(10')-triene-7' α -yl) undecanamide (16, X = Cl)

To diacetate amide 15, dissolved in 5 ml of acetone, was added a solution of sodium acetate (2.6 equivalents) in acetic acid and water (1:11.3 v/v) and then, was treated with tertbutyl hypochlorite (1 eq.) prepared from *t*-butanol (4 ml) and Javel water (Javex 6.1%, 50 ml). The clear solution was warmed to 55°C and stirred for 1 h. Afterwards, the solvent was evaporated to dryness. The residue was dissolved in ether (100 ml) and water was added (20 ml). The organic phase was washed with water, dried with anhydrous $MgSO_4$ and evaporated to dryness. The residue was purified by chromatography on silica gel carried out with mixture of EtOAc/hexane (3:7 v/v) to give the *N*-butyl, *N*-methyl-11-(16' α -chloro-3'-acetoxy-17'-oxo-estra-1',3',4'-(10')-triene-7' α -yl) undecanamide 16, X=Cl (115 mg, 89%) as colorless oil; 1H -NMR ν ($CDCl_3$) 0.92 and 0.95 (2t, 3H, $J=7.0$ Hz, $N(CH_3)_2CH_3$), 0.96 (s, 3H, 18'- CH_3), 2.28 (s, 3H, 3'- $OCOCH_3$), 2.80 app (d, 1H, $J=16.6$ Hz, part of ABX system, 6'-H) 2.90 and 2.96 (2s, 3H, $N-CH_3$), 3.24 and 3.35 (2t_{app}, 2H, $J=7.4$ Hz, $-N-CH_2-$), 4.46 (d, 1H, $J=6.6$ Hz, 16' β -H), 6.82 (broad s, 1H, 4'-H), 6.86 (dd, 1H, $J=9.1$ Hz and $J_2=2.6$ Hz, 2'-H), 7.29 (d, 1H, $J=9.1$ Hz, 1'-H); IR ν_{max} (neat) 1750, 1640, 1205 cm^{-1} ; MS m/e 601, 599 (M^+ , 24%, 68%), 142 ($C_8H_8CON(CH_3)C_8H_8$, 100%), 114 ($C_8H_8(CH_3)NCO$, 93%).

N-butyl, N-methyl-11-(16 α -chloro-3',17'-dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 139") and ("EM 170")

A stirred solution of haloketone amide (16, X=Cl) in anhydrous tetrahydrofuran (THF) (10 ml) under argon was chilled to -70°C with 2-propanol/dry ice bath. A solution of 1.0 M of lithium aluminium hydride (2 eq.) was then added dropwise. After 30 min, the reaction was allowed to return slowly at 0°C for 5 min, then was quenched by the dropwise addition of a mixture of THF-EtOAc (5 ml) (1:1 v/v) and acidified at pH ~ 4 with (10%) HCl. The mixture was stirring for 5 min at room temperature and then extracted with EtOAc. The organic phase was washed with water, dried on anhydrous Na₂SO₄ and evaporated under reduced pressure. The residue was chromatographed on silica gel with a mixture of EtOAc/hexane (4:6 v/v) as eluant:

N-butyl, N-methyl-11-(16' α -chloro-3'17' α -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 170")

(15 mg, 29%) as colorless oil; analytical sample was obtained by HPLC purification; ¹H-NMR δ (CDCl₃, 400 MHz) 0.79 (s, 3H, 18'-CH₃), 0.93 and 0.96 (2t, 3H, J=7.3 Hz, N(CH₃), CH₃), 2.80 (2H, J_{1,2}=17.1 Hz and J_{2,3}=4.5 Hz, $\Delta\delta$ =24.34 (Hz, system ABX, 6'-H), 2.94 and 2.99 (2s, 3H, N-CH₃), 3.26 (dd, J₁=7.6 Hz and J₂=7.4 Hz) and 3.32-3.43 (m)-[2H, -N-CH₂-], 3.71 (d, 1H, J=4.5 Hz, 17' β -H), 4.63 (ddd, 1H, J_{1,2}=10.2 Hz, J_{1,3}=4.5 Hz and J_{2,3}=3.9 Hz, 16' β -H), 6.50 (d, 1H, J=24 Hz, 3'-OH), 6.60 (d, 1H, J=2.5 Hz, 4'-H), 6.66 (dd, 1H, J₁=8.4 Hz and J₂=2.5 Hz, 2'-H), 7.14 (d, 1H, J=8.5 Hz, 1'-H); IR ν_{\max} (neat) 3300, 1615, 1495 cm⁻¹; MS m/e

SUBSTITUTE SHEET


561,559 (M^+ , 40%, 100%), 523 (M^+-HCl , 20%), 142 ($C_2H_5CON(CH_3)C_6H_5^+$, 44%), 114 ($C_6H_5(CH_3)NCO^+$, 37%); Exact mass calculated for $C_{24}H_{34}O_3N^+Cl$ 559.3785, found 559.3821;

- and -

-N-butyl, N-methyl-11-(16' α -chloro-3',17' β -dihydroxy-estra-1'3',5'(10')-trien-7' α -yl) undecanamide ("EM 139")

(25 mg, 55%) as a colorless oil; analytical sample was obtained by HPLC purification: 1H -NMR δ ($CDCl_3$, 400 MHz), 0.81 (s, 3H, 18'- CH_3), 0.93 and 0.96 (2t, 3H, $J=7.3$ Hz, $(CH_2)_3CH_3$), 2.78 (2H, $J_{6,7}=16.2$ Hz and $J_{6,8}=4.5$ Hz, $\Delta\delta=24.34$ Hz, system ABX, 6'-H), 2.94 and 2.99 (2s, 3H, N- CH_3), 3.27 (dd, $J_1=7.6$ Hz and $J_2=7.5$ Hz) and 3.31-3.45 (m) [2H, -N- CH_2 -], 3.86 (dd, 1H, $J_{1,2}=3.4$ Hz and $J_{1,3}=5.9$ Hz, 17' α -H), 4.11 (ddd, 1H, $J_{1,2}=10.8$ Hz, $J_{1,3}=5.9$ Hz and 4.11 (ddd, 1H, $J_{1,2}=10.8$ Hz, $J_{1,3}=5.9$ Hz and $J_{1,4}=2.5$ Hz, 16' β -H), 6.56 (d, 1H, $J=19.7$ Hz, 3'-OH), 6.61 (d, 1H, $J=2.5$ Hz, 4'-H), 6.66 (dd, 1H, $J_1=8.4$ Hz and $J_2=2.6$ Hz, 2'-H), 7.13 (d, 1H, $J=8.4$ Hz, 1'-H); IR ν_{max} (neat) 3320, 1615, 1490 cm^{-1} ; MS m/e 561,559 (M^+ , 38%, 100%), 523 (M^+-HCl , 16%), 142 ($C_2H_5CON(CH_3)C_6H_5^+$, 80%), 114 ($C_6H_5(CH_3)NCO^+$, 76%); exact mass calculated for $C_{24}H_{34}O_3N^+Cl$ 559.3785, found 559.3825.

Schema 3



SUBSTITUTE SHEET

N-*n*-butyl, *N*-methyl-11-(16' α -bromo-3'-acetoxy-17'-~~oxo~~-estra-1',3',5'-(10'),trien-7' α -yl) undecanamide (16, X=Br)

To the above diacetate 15 (244 mg, 0.40 mmol) dissolved in 10 ml of acetic acid was added dropwise with stirring within 10 minutes and at room temperature, a brominating solution composed of 50 mg (0.6 mmol) of sodium acetate, 1.6 ml of acetic acid, 0.04 ml of water and 63.9 mg (0.02 ml, 0.40 mmol) of bromine. During the course of this reaction, a red coloration appeared and disappeared. To the solution, 50 ml of ether was added and the organic phase was washed with water (4 x 50 ml) followed by a saturated sodium bicarbonate solution (2 x 50 ml) and finally with water (3 x 50 ml). The combined phase was dried over anhydrous magnesium sulfate and the solvent was removed in vacuo. The residue was chromatographed on silica gel (Kieselgel, 60F254, Merck, 0.063-0.200 mm). Elution with a mixture of hexane-ethyl acetate (4:1 v/v) yielded *N*-butyl, *N*-methyl-11-(16 α -bromo-3'-acetoxy-17'-oxo-estra-1',3',5'(10'),trien-7'- α -yl) undecanamide (16, X=Br) (201 mg, 78%) as colorless oil (201 mg, 78%), as colorless oil; $^1\text{H-NMR}$ ϕ (CDCl_3), 0.94 (s, 3H, 18'-CH₃), 2.28 (s, 3H, 3'-OCOCH₃), 2.82 app (d, 1H, J=16.4 Hz, part of ABX system, 6'-H), 2.90 and 2.96 (2s, 3H, N-CH₃), 3.24 and 3.35 (2t_{app}, 2H, J=7.7 Hz, -N-CH₂-), 4.58 (t, 1H, J=3.6 Hz, 16 β -H), 6.82 (broad s, 1H, 4'-H), 6.88 (dd, 1H, J=8.0 Hz and J₂= 4.0 Hz, 2'-H), 7.29 (d, 1H, J=8.0 Hz, 1'-H); MS m/e 644 (M^+ , 7%), 565 ($\text{M}^+ - \text{Br}$, 77%), 522 ($\text{M}^+ - \text{Br} - \text{COCH}_3$, 55%), 142 ($\text{C}_2\text{H}_5\text{CON}(\text{CH}_3)\text{C}_4\text{H}_9^+$, 67%), 114 ($\text{C}_4\text{H}_9(\text{CH}_3)\text{NCO}^+$, 66%), 88 (100%).

N-butyl, N-methyl-11-(16' α -bromo-3',17'-dihydroxy-estra-1',3,4'(10')-trien-7' α -yl) undecanamide ("EM 105") and ("EM 171")

A solution of bromoketone amide 16 (X=Br) (295 mg, 0.46 mmol) in anhydrous tetrahydrofuran (10 ml) under argon was chilled to -70°C and a solution of 1.0 M of lithium aluminium hydride in ether (0.92 ml, 0.92 mmol) was added dropwise with rapid magnetic stirring. After 30 min, the reaction was quenched by the dropwise addition of a mixture of THF-ethyl acetate (1:1 v/v) and acidified by 10% hydrochloric acid. The mixture was stirring for 5 min at room temperature and then extracted with ethyl acetate. The organic phase was washed with water, dried on anhydrous sodium sulfate and evaporated to dryness under reduced pressure. The residue was purified by chromatography on silica gel. Elution with a mixture of hexane-ethyl acetate (7:3 v/v) gave:

N-n-butyl, N-methyl-11-(16' α -bromo-3',17' α -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 171")

(63 mg, 21%) as colorless oil; $^1\text{H-NMR}$ δ (CDCl_3 , 400 MHz) 0.81 (s, 3H, 18'-CH₃), 0.93 and 0.96 (2t, 3H, $J=7.3$ Hz, N(CH₂), CH₂), 2.79 (2H, $J_{6,6}=16.6$ Hz, $J_{6,7}=4.7$ Hz, $=\Delta\delta=24.34$ Hz, system ABX, 6'-H), 2.94 and 2.99 (2s, 3H, N-CH₃), 3.27 (dd, 2H, $J_1=7.7$ Hz and $J_2=7.5$ Hz, -N-CH₂-), 3.31-3.44 (m, 2H, -N-CH₂-), 3.66 (dd, 1H, $J_{17,17}=1.4$ Hz, $J_{17,16}=4.3$ Hz, 17' β -H), 4.68 (dt, 1H, $J_{16,17}=4.3$ Hz, m, $J_{16,15}=9.7$ Hz, 16' β -H), 6.60 (d, 1H, $J=2.4$ Hz, 4'-H), 6.65 (dd, 1H, $J=8.5$ Hz and $J_2=2.5$ Hz, 2'-H), 7.14 (d, 1H, $J=8.5$ Hz, 1'-H); IR ν_{max} (neat) 3300, 1615, 1495 cm^{-1} ; MS m/e 605, 603 (M^+ , 17%), 523 (M^+-HBr , 81%), 142 ($\text{C}_8\text{H}_8\text{CON}(\text{CH}_3)\text{C}_4\text{H}_9^+$, 100%), 114 ($\text{C}_8\text{H}_8(\text{CH}_3)\text{NCO}^+$, 97%); Exact mass calculated for $\text{C}_{31}\text{H}_{48}\text{O}_2\text{N}^+\text{Br}$ 603.8289, found 603.3304.

SUBSTITUTE SHEET

and

~~N-n-butyl, N-methyl-11-(16' α -bromo-3',17' β -dihydroxy-estra-1',3',-~~
5'(10')-trien-7 α -yl) undecanamide ("EM 105")

(170 mg, 50%) as a colorless oil; analytical sample was obtained by HPLC purification; $^1\text{H-NMR}$ δ (CDCl_3 , 400 MHz), 0.80 (s, 3H, 18, - CH_3), 0.93 and 0.96 (2t, 3H, $J=7.3$ Hz, $\text{N}(\text{CH}_2)_2\text{CH}_3$), 2.80 (2H, $J_{6,6}=16.4$, $J_{6,7}=4.6$ Hz, $\Delta\delta=24.34$ Hz, system ABX, 6'-H), 2.94 and 2.99 (2s, 3H, N-CH_3), 3.27 (dd, 2H, $J_1=7.7$ Hz and $J_2=7.5$ Hz, - N-CH_2 -), 3.31-3.45 (m, 2H, - N-CH_2 -), 4.02 (dd, 1H, $J_{17,17}=3.7$ Hz, and $J_{17,16}=6.1$ Hz, 17' α -H), 4.15 (ddd, 1H, $J_{16,15}=10.2$ Hz, $J_{16,17}=6.1$ Hz and $J_{16,15}=2.9$ Hz, 16' β -H), 6.61 (d, 1H, $J=2.5$ Hz, 4'-H), 6.66 (dd, 1H, $J=8.4$ Hz and $J_2=2.5$ Hz, 2'-H), 7.12 (d, 1H, $J=8.4$ Hz, 1'-H); IR ν_{max} (neat) 3320, 1610, 1490 cm^{-1} ; MS m/e 605, 603 (M^+ , 29%), 523 (M^+-HBr , 100%), 142 ($\text{C}_2\text{H}_5\text{CON}(\text{CH}_2)_2\text{C}_2\text{H}_5^+$, 70%), 114 ($\text{C}_2\text{H}_5(\text{CH}_2)_2\text{NCO}^+$, 60%); Exact mass calculated for $\text{C}_{34}\text{H}_{50}\text{O}_2\text{N}^+\text{Br}$ 603.3289, found 603.3289.

Antiestrogens useful in the combination therapy of the invention include but are not limited to Tamoxifen, commercially available from Imperial Chemical Industries, and EM 139, EM 170 and EM 171 whose synthesis are set forth above. Some steroidal antagonists also function as inhibitors of sex steroid formation. The antiestrogens EM 139, EM 170 and EM 171, for example, exhibit the dual function of acting as sex steroid formation inhibitors. For this reason, a combination therapy requiring both an inhibitor of sex steroid formation and a steroidal antagonist may be produced by administering a single active compound (alone or together with diluent) capable of performing both functions. Another example of a dual function active ingredient is the antiandrogen EM 101 which has also shown an inhibitory effect on sex steroid formation.

The inhibitor of sex steroid biosynthesis is preferably capable of acting at least in peripheral tissues (extra-testicular and extra-adrenal). In preferred embodiments, it is used in association with an antiandrogen, and with an LHRH agonist or LHRH antagonist. The use of an LHRH agonist is the more preferred method of chemical castration. Surgical castration may alternatively be used as a means of inhibiting testicular hormonal secretions, but chemical castration is preferred.

By the term "LHRH agonist" is meant synthetic analogues of the natural luteinizing hormone-releasing hormone (LHRH), a decapeptide of the structure: L-pyroglutamyl-L-histidyl-L-tryptophyl-L-seryl-L-tyrosyl-glycyl-L-leucyl-arginyl-L-prolylglycyl-NH₂.

Typical suitable LHRH agonists include nonapeptides and decapeptides represented by the formula: L-pyroglutamyl-L-histidyl-L-tryptophyl-L-seryl-L-tyrosyl-X-Y-L-arginyl-L-prolyl-Z wherein X is D-tryptophyl, D-leucyl, D-alanyl, iminobenzyl-D-histidyl, 3-(2-naphthyl)-D-alanyl, O-terbutyl-D-seryl, D-tyrosyl, D-lysyl, D-phenylalanyl or N-methyl-D-alanyl and Y is L-leucyl, D-leucyl, N^α-methyl-D-leucyl, N^α-methyl-L-leucyl or D-alanyl and wherein Z is glycyl-NHR₁ or NHR₁, wherein R₁ is H, lower alkyl or lower haloalkyl. Lower alkyl includes, for example, methyl.

SUBSTITUTE SHEET

Haloloweralkyl includes, for example, $-\text{CF}_2-$, $-\text{CH}_2\text{CF}_2-$, $-\text{CF}_2\text{CH}_2-$, and the like. Fluorine is a preferred halogen.

Preferred nonapeptides wherein Y is L-leucyl and X is an optically active D-form of selected amino acids and Z is NHC_2H_5 , are [D-Trp⁶, des-Gly-NH₂¹⁰]LHRH ethylamide (X=D-Trp⁶); [D-Ser(t-BuO)⁶, des-Gly-NH₂¹⁰]LHRH ethylamide [X=D-Ser(t-BuO⁶)]; [D-Leu⁶, des-Gly-NH₂¹⁰]LHRH ethylamide (X=D-Leu⁶, [D-His(Bzl)⁶, des-Gly-NH₂¹⁰]LHRH ethylamide (X=iminobenzyl-D-His⁶) and [D-Ala⁶, des-Gly-NH₂¹⁰]LHRH ethylamide (X=D-Ala⁶).

Preferred decapeptides include [D-Trp⁶]LHRH wherein X=D-Trp, Y=L-leucyl, Z=glycyl-NH₂, [D-Phe⁶]LHRH wherein X=D-phenylalanyl, Y=L-leucyl and Z=glycyl-NH₂, or [D-Nal(2)⁶]LHRH which is [3-(2-naphthyl)-D-Ala⁶]LHRH wherein X=3-(2-naphthyl)-D-alanyl, Y=L-leucyl and Z=glycyl-NH₂.

Other LHRH agonists useful within the scope of this invention are the α -aza analogues of the natural LHRH, especially, [D-Phe⁶, Azgly¹⁰]LHRH, [D-Tyr(-Me)⁶, Azgly¹⁰]LHRH, and [D-Ser(t-BuO)⁶, Azgly¹⁰]LHRH disclosed by A.S. Dutta et al. in J. Med. Chem., 21, 1018 (1978) and U.S. Pat. No. 4,100,274 as well as those disclosed in U.S. Pat. Nos. 4,024,248 and 4,118,483.

Typical suitable LHRH antagonists include [N-Ac-D-p-Cl-Phe^{1,2}, D-Phe³, D-Arg⁶, D-Ala¹⁰]LHRH disclosed by J. Ercheggi et al., Biochem. Biophys. Res. Commun. 100, 915-920 (1981); [N-Ac-D-p-Cl-Phe^{1,2}, D-Trp³, D-Arg⁶, D-Ala¹⁰]LHRH disclosed by D.H. Coy et al., Endocrinology, 110: 1445-1447 (1982); [N-Ac-D-(3-(2-naphthyl)-OAla)¹, D-p-Cl-Phe², D-Trp³, D-hArg(Et)⁶, D-Ala¹⁰]LHRH and [N-Ac-Pro¹, D-pF-Phe², D-(3-(2-naphthyl)-Ala³,⁶)]LHRH disclosed by J.J. Nestor et al. J. Steroid Biochem., 20 9no. 6B, 1366 (1984); the nona- and decapeptide analogs of LHRH useful as LHRH antagonists disclosed in U.S. Pat. No. 4,481,190 (J.J. Nestor et al.); analogs of the highly constrained cyclic antagonist, cycle [A¹

Pro¹, D-p-Cl-Phe², D-Trp^{3,4}, N-Me-Leu⁵, β-Ala¹⁰]-LHRH disclosed by J. Rivier, J. Steroid Biochem., 20 (no. 6B), 1365 (1984); and [N-Ac-D-(3-(2-naphthyl)-Ala¹, D-p-F-Phe², D-Trp³, D-Arg⁴]-LHRH disclosed by A. Corbin et al., J. Steroid Biochem. 20 (no. 6B) 1369 (1984).

Preferred nonapeptides wherein Y is L-leucyl and X is an optically active D-form of selected amino acids and Z is NHC₂H₅, are [D-Trp⁶, des-Gly-NH₂¹⁰]-LHRH ethylamide (X=D-Trp⁶); [D-Ser-(t-BuO)⁶, des-Gly-NH₂¹⁰]-LHRH ethylamide [X=D-Ser-(t-BuO)⁶]; [D-Leu⁶, des-Gly-NH₂¹⁰]-LHRH ethylamide (X=D-Leu⁶, [D-His(Bzl)⁶, des-Gly-NH₂¹⁰]-LHRH ethylamide (X=iminobenzyl-D-His⁶) and [D-Ala⁶, des-Gly-NH₂¹⁰]-LHRH ethylamide (X=D-Ala⁶).

Preferred decapeptides include [D-Trp⁶]-LHRH wherein X=D-Trp, Y=L-leucyl, Z=glycyl-NH₂, [D-Phe⁶]-LHRH wherein X=D-phenylalanyl, Y=L-leucyl and Z=glycyl-NH₂ or [D-Nal(2)⁶]-LHRH which is [3-(2-naphthyl)-D-Ala⁶]-LHRH wherein X=3-(2-naphthyl)-D-alanyl, Y=L-leucyl and Z=glycyl-NH₂.

Other LHRH agonists useful within the scope of this invention are the α-aza analogues of the natural LHRH, especially, [D-Phe⁶, Azgly¹⁰]-LHRH, [D-Tyr(-Me)⁶, Azgly¹⁰]-LHRH, and [D-Ser-(t-BuO)⁶, Azgly¹⁰]-LHRH disclosed by A.S. Dutta et al. in J. Med. Chem., 21, 1018 (1978) and U.S. Pat. No. 4,100,274 as well as those disclosed in U.S. Pat. Nos. 4,024,248 and 4,118,483.

Typical suitable LHRH antagonists include [N-Ac-D-p-Cl-Phe^{1,2}, D-Phe³, D-Arg⁴, D-Ala¹⁰]-LHRH disclosed by J. Ercheggi et al., Biochem. Biophys. Res. Commun. 100, 915-920 (1981); [N-Ac-D-p-Cl-Phe^{1,2}, D-Trp³, D-Arg⁴, D-Ala¹⁰]-LHRH disclosed by D.H. Coy et al., Endocrinology, 110: 1445-1447 (1982); [N-Ac-D-(3-(2-naphthyl)-OAla)¹, D-p-Cl-Phe², D-Trp³, D-hArg(Et)⁴, D-Ala¹⁰]-LHRH and [N-Ac-Pro¹, D-pF-Phe², D-(3-(2-naphthyl)-Ala³,⁶]-LHRH disclosed by J.J. Nestor et al. J. Steroid Biochem., 20 9no. 6B), 1366 (1984); the nona- and decapeptide analogs of LHRH useful as

SUBSTITUTE SHEET

LHRH antagonists disclosed in U.S. Pat. No. 4,481,190 (J.J. Nestor et al.); analogs of the highly constrained cyclic antagonist, cycle [Δ^1 Pro¹, D-p-Cl-Phe², D-Trp³,⁴, N-Me-Leu⁵, β -Ala¹⁰]-LHRH disclosed by J. Rivier, J. Steroid Biochem., 20 (no. 6B), 1365 (1984); and [N-Ac-D-(3-(2-naphthpyl)-Ala¹, D-p-F-Phe², D-Trp³, D-Arg⁶)]LHRH disclosed by A. Corbin et al., J. Steroid Biochem. 20 (no. 6B) 1369 (1984).

Other LHRH agonist and antagonist analogs are disclosed in LHRH and its Analogs (B.H. Vickery et al. eds, at pages 3-10 (J.J. Nestor), 11-22 (J. Rivier et al.) and 22-33 (J.J. Nestor et al.) as well as in The Case for LHRH agonists (Clinical Oncology, Furr and Denis, eds), Baillière Tindall, vol. 2, no. 3, pp. 559-570, 1988).

The LHRH agonists and antagonists useful in this invention may conveniently be prepared by the method described by Stewart et al. in "Solid Phase Peptide Synthesis" (published in 1969 by Freeman & Co., San Francisco, page 1) but solution phase synthesis may also be used.

The nona- and decapeptides used in this invention are conveniently assembled on a solid resin support, such as 1% cross-linked Pro-merri-field resin by use of an automatic peptide synthesizer. Typically, side-chain protecting groups, well known to those in the peptide arts, are used during the dicyclohexylcarbodiimidecatalyzed coupling of a tert-butyloxycarbonylamino acid to the growing peptide attached to a benzhydryl-amide resin. The tert-butyloxycarbonyl protecting groups are removed at each stage with trifluoroacetic acid. The nona- or decapeptide is cleaved from the resin and deprotected by use of HF. The crude peptide is purified by the usual techniques, e.g., gel filtration and partition chromatography and optionally lyophilization. See also D.H. Coy et al., J. Med. Chem. 19, pages 423-425 (1976).

In this invention, the LHRH agonist or antagonist, the 5 α -reductase inhibitor, the antiandrogen, the antiestrogen, and, where applicable, the

SUBSTITUTE SHEET

inhibitor of 3β - and 17β -hydroxysteroid dehydrogenase activities are administered as pharmaceutical compositions via topical, parenteral or oral means. The LHRH agonist or antagonist is administered parenterally, i.e., intramuscularly, subcutaneously or intravenously by injection or infusion by nasal drops or by suppository. The LHRH agonist or antagonist may also be microencapsulated in or attached to a biocompatible, biodegradable polymer, e.g., poly(d,l-lactide-co-glycolide) and subcutaneously or intramuscularly injected by a technique called subcutaneous or intramuscular depot to provide continuous, slow release of the LHRH agonist or antagonist over a period of 30 days or longer. The most preferred route of administration of the LHRH agonist or antagonist is subcutaneous or intramuscular depot injection. Preferably the antiestrogen will be administered orally. Preferably, the 5α -reductase inhibitor, the antiandrogen, the antiestrogen, the inhibitor of 3β -HSD and the inhibitor of 17β -HSD can also be administered orally. The antiestrogen, an inhibitor of 3β -HSD and inhibitor of 17β -HSD can also be administered in a slow release formulation, e.g. poly(d,l-lactide-coglycolide) or as implants.

The amount of each component administered is determined by the attending clinicians taking into consideration the etiology and severity of the disease, the patient's condition and age, the potency of each component and other factors. According to this invention, the following dosage ranges are suitable.

The LHRH agonist or antagonist is generally administered at from about 10 to 5000 μg per day with contemplated dosage ranges of about 10 to 1500 μg per day and about 250 (preferably 50 μg to 500 μg per day) for the LHRH agonist and to about 100 to 2000 μg per day for the LHRH antagonist being preferred.

SUBSTITUTE SHEET

In the most preferred embodiment of this invention, the LHRH agonist or antagonist is administered subcutaneously in a daily dose of 500 μg for the first 30 days and thereafter subcutaneously in a daily dose of 250 μg regardless of the patients' body weight. When the LHRH agonist or antagonist is administered, once every 30-day period is used, with a dose of 750 to 15,000 μg per 30-day period being preferred. Similar daily delivery doses are used for longer-term controlled release formulations.

The inhibitors of 3β -HSD and 17β -HSD are preferably administered in dosages ranging from about 0.1 to 25 mg/kg per day with 200 mg per day in two equally divided doses being preferred.

The antiestrogen compositions are administered in a dosage range of about 0.05 to 25 mg/kg body weight per day, with 20 mg, especially 40 mg, in two equally divided doses being preferred.

The 5α -reductase inhibitor compositions are administered in a dosage ranging from 0.1 to 25 mg/kg per day with 50 mg per day in two equivalent doses being preferred.

The antiandrogen and aromatase inhibitor compositions are administered in a dosage range of 0.5 to 25 mg/kg body weight per day with 750 mg per day in three equally divided doses being preferred.

The LHRH agonist or antagonist, antiestrogen, antiandrogen, an inhibitor of aromatase, 17β -HSD and 3β -HSD each may be administered separately or when the modes of administration are the same, all or at least two of them may be administered in the same composition, but in any case the preferred ratio of LHRH agonist to antiestrogen, to antiandrogen to inhibitor of 17β -HSD and administered daily will be about 250 μg of LHRH agonist to about 750 mg of antiandrogen, about 40 mg of antiestrogen, to about 40 mg of inhibitor of 17β -HSD and about 40 mg of inhibitor of 3β -HSD.

In the therapy of prostate cancer, combining the administration of an LHRH agonist or antagonist, an antiestrogen, an antiandrogen and an inhibitor of 17β -HSD, the dosages preferable are as follows: the LHRH

SUBSTITUTE SHEET

agonist or antagonist is generally administered at from about 10 to 2000 μg per day, with contemplated dosage ranges of 10 to 500 μg per day, 50-250 μg per day and 250 to 500 μg per day being preferred. In the most preferred embodiment of this aspect of this invention, the LHRH agonist or antagonist is administered subcutaneously in a daily dose of 500 μg for the first 30 days and thereafter subcutaneously in a daily dose of 250 μg regardless of the patients' body weight. When the LHRH agonist or antagonist is administered, once every 30-day period, by intramuscular or subcutaneous depot injection, a dose from about 300 to 60000 (occasionally 10000) μg per 30-day period is used, with a dose of 750 to 2000 μg per 30-day period being preferred. The antiandrogen composition is generally administered in a dosage range of about 0.5 to 25 mg/kg (body weight) per day with 400 especially 750 mg per day in three equally divided doses being preferred. The antiestrogen and inhibitor of $17\beta\text{-HSD}$ and $3\beta\text{-HSD}$ activities are administered in a dosage range of about 0.1 to 25 mg/kg body weight per day, with 100 mg in two, preferably with 50 mg in two, equally divided doses being preferred.

The LHRH agonist or antagonist, antiandrogen, antiestrogen, $5\alpha\text{-reductase}$ inhibitor, inhibitor of $17\beta\text{-HSD}$, inhibitor of $3\beta\text{-HSD}$, inhibitor of aromatase, each may be administered separately or when the modes of administration are the same, all or two or three of them may be administered in the same composition, but in any case the preferred ratio of LHRH agonist to antiandrogen to antiestrogen administered daily will be about 750 μg of LHRH agonist to about 250 mg of antiandrogen to preferably 40 mg of antiestrogen.

In the therapy of prostate cancer, according to this invention, it is preferred that the LHRH agonist is $[\text{D-Trp}^6, \text{des-Gly-NH}_2^{10}] \text{LHRH ethylamide}$ be administered subcutaneously in single daily dose of 500 μg for the first thirty (30) days of treatment and thereafter in a single daily dose of 250 μg .

SUBSTITUTE SHEET

In the combination therapy of prostate cancer according to this invention, the administration of the antiandrogen, antiestrogen, inhibitor of 17β -HSD, inhibitor of 5α -reductase, inhibitor of aromatase, and inhibitor of 3β -HSD, LHRH agonist or LHRH antagonist can be started in any order of sequence. Preferably, the administration of the antiandrogen and 5α -reductase inhibitor, are started before (preferably two to four hours before) the administration of the LHRH agonist or LHRH antagonist is started. Orchiectomy can replace LHRH agonist or antagonist. Preferably, the administration of the inhibitor of 17β -HSD and inhibitor of 3β -HSD is started on the same day as the administration of the LHRH agonist or LHRH antagonist. However, the attending clinician may elect to start administration of the LHRH agonist or antagonist simultaneously with the antiandrogen, antiestrogen inhibitor of 17β -HSD and inhibitor of 3β -HSD.

When patients whose testes have already been surgically removed are treated according to this invention, the administration and dosage of the antiandrogen and the other components of the therapy (except the LHRH agonist or antagonist which is not used) are the same as indicated for the therapy in which the LHRH agonist or antagonist is used.

The LHRH agonists or antagonists useful in the present invention are typically amorphous solids which are freely soluble in water or dilute acids, e.g., HCl, H_2SO_4 , citric, acetic, mandelic or fumaric. The LHRH agonist or antagonist for subcutaneous injection is supplied in vials containing 5 ml of sterile solution with the LHRH agonist or antagonist at a concentration of about 1.0 mg/ml.

A typical pharmaceutical composition of the LHRH agonist or antagonist includes the LHRH agonist or antagonist or a pharmaceutically acceptable acid salt thereof, benzyl alcohol, a phosphate buffer (pH 6.0-6.5) and sterile water.

The LHRH agonist or antagonist for intramuscular or subcutaneous depot injection may be microencapsulated in a biocompatible, biodegradable polymer, e.g., poly (d,l-lactide-co-glycolide) by a phase separation process or formed into a pellet. The microspheres may then be suspended in a carrier to provide an injectable preparation or the depot may be injected in the form of a pellet. See also European patent application EPA No. 58,481 published Aug. 25, 1982 for solid compositions for subdermal injection or implantation or liquid formulations for intramuscular or subcutaneous injections containing biocompatible, biodegradable polymers such as lactide-glycolide copolymer and an LHRH agonist, e.g. D-Ser-t-BuO⁶, Azgly¹⁰-LHRH. These formulations permit controlled release of the peptide.

The inhibitors of 17 β -HSD, 3 β -HSD, aromatase and 5 α -reductase are typically compounded in customary ways for oral administration, e.g., in tablets, capsules and the like. These compounds useful in the present invention are typically formulated with conventional pharmaceutical excipients, e.g., spray dried lactose and magnesium stearate into tablets or capsules for oral administration. The antiestrogens, when used with the invention, are typically compounded in customary ways for oral administration, e.g., in capsules, tablets, as dragees or even in liquid form, e.g., suspensions or syrups. One or more of the active substances, with or without additional types of active agents, can be worked into tablets or dragee cores by being mixed with solid, pulverulent carrier substances, such as sodium citrate, calcium carbonate or dicalcium phosphate, and binders such as polyvinyl pyrrolidone, gelatin or cellulose derivatives, possibly by adding also lubricants such as magnesium stearate, sodium lauryl sulfate, "Carbowax" or polyethylene glycols. Of course, taste-improving substances can be added in the case of oral administration forms.

SUBSTITUTE SHEET

The therapeutically active antiestrogen compound should be present in a concentration of about 0.5-90% by weight of the total mixture, i.e., in amounts that are sufficient for maintaining the above-mentioned dosage range.

As further forms, one can use plug capsules, e.g., of hard gelatin, as well as closed soft-gelatin capsules comprising a softener or plasticizer, e.g. glycerine. The plug capsules contain the active substance preferably in the form of granulate, e.g., in mixture with fillers, such as lactose, saccharose, mannitol, starches, such as potato starch or amylopectin, cellulose derivatives or highly-dispersed silicic acids. In soft-gelatin capsules, the active substance is preferably dissolved or suspended in suitable liquids, such as vegetable oils or liquid polyethylene glycols.


In place of oral administration, the active compounds may be administered parenterally. In such case, one can use a solution of the active substance, e.g., in sesame oil or olive oil. One or more of the active substances (antiestrogen or inhibitor of 17β -HSD and 3β -HSD can be microencapsulated in or attached to a biocompatible, biodegradable polymer, e.g. poly(d,l-lactide-co-glycolide) and subcutaneously or intramuscularly injected by a technique called subcutaneous or intramuscular depot to provide continuous slow release of the compound(s) for a period of 2 weeks or longer.

In the most preferred aspect of this invention, the LHRH agonist is [D-Trp⁶,des-Gly-NH₂,¹⁰] LHRH ethylamide which is administered subcutaneously in single daily dose of 500 µg for the first thirty (30) days of treatment and thereafter in a single daily dose of 250 µg; the antiandrogen is EM 101 which is administered orally in three equally divided daily doses of 250 mg; and the inhibitor of sex steroid biosynthesis is EM 139 and/or MK 906 administered orally in two equally divided doses of 50 mg every 12 hours.

The inhibitor(s) of sex steroid biosynthesis and the antiandrogen are preferably administered to a male in need of the prostate cancer treatment of this invention two to four hours before the LHRH agonist or antagonist is administered, but the attending clinician may elect to start administration of the LHRH agonist or antagonist, the antiandrogen and the inhibitor of steroid biosynthesis simultaneously. When the antiandrogen and sex steroid inhibitor are particularly effective, both chemical (LHRH agonist or antagonist) and surgical castration may be avoided. Especially, when patients whose testes have already been surgically removed are treated according to this invention, no LHRH agonist or antagonist need to be used but other dosages remain the same.

The terms and descriptions used herein are preferred embodiments set forth by way of illustration only, and are not intended as limitations on the many variations which those of skill in the art will recognize to be possible in practicing the present invention as defined by the following claims.

1. A method for treating prostate cancer in humans or other warm-blooded animals in need of such treatment, said method comprising the steps of blocking androgen receptors by administering a therapeutically effective amount of an antiandrogen having, as part of its molecular structure, a substituted or unsubstituted androgenic nucleus of the formula:



and having as another part of its molecular structure, at least one side chain represented by the formula:

$-R^1[-B-R^2-]_x L-G$ said chain being substituted onto said androgenic nucleus at a position selected from the group consisting of 6, 7, 14, 15, 16 and 17, wherein:

x is an integer from 0 to 6, wherein at least one of L and G is a polar moiety distanced from said ring carbon by at least three intervening atoms, and wherein:

R^1 and R^2 are independently either absent or selected from the group consisting of straight- or branched-chain alkylene, straight- or branched chain alkynylene, straight- or branched-chain alkenylene, phenylene and fluoro-substituted analogs of the foregoing;

B is either absent or selected from the group consisting of $-O-$, $-S-$, $-Se-$, $-SO-$, $-SO_2-$, $-NR^3-$, $-SiR^3_2-$, $-CR^3OR^3-$, $-NR^3CO-$, $-NR^3CS-$, $-CONR^3-$, $-CSNR^3-$, $-COO-$, $-COS-$, $-SCO-$, $-CSS-$, $-SCS-$, $-OCO-$ and phenylene (R^3 being hydrogen or lower alkyl);

L is either a moiety which together with G , forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of lower alkyl, $-CONR^4-$, $-CSNR^4-$, $-NR^4CO-$, $-NR^4CS-$, $-NR^4CONR^4-$

$\begin{matrix} NR^4 \\ || \\ -NR^4C-NR^4- \end{matrix}$, $-SO_2NR^4-$, $-CSS-$, $-SCS-$, $-(NO)R^4-$, $-(PO)R^4-$, $-NR^4COO-$,

SUBSTITUTE SHEET

-NR³SO₂-, -O-, -NR⁴-, -S-, -SO- and -SO₂- (R⁴ and R⁵ being independently selected from the group consisting of hydrogen and lower alkyl; and R⁶ being selected from the group consisting of hydrogen, nitrile and nitro); and

G is either a moiety which together with L forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, (C₃-C₆)cycloalkyl, bromo(lower)alkyl, chloro(lower)alkyl, fluoro(lower)alkyl, cyano(lower)alkyl, carboxy(lower)alkyl, (lower)alkoxycarbonyl(lower)alkyl, (C₆-C₁₀)aryl, (C₆-C₁₁)arylalkyl, di(lower)alkylamino(lower)alkyl, fluoro-substituted analogs of the foregoing and a moiety which together with L forms a heterocyclic ring having at least one nitrogen atom;

Said method of treatment further comprising the step of inhibiting sex steroid formation by administering therapeutically effective amount of at least one sex steroid formation inhibitor.

2. The method of Claim 1 wherein said inhibition of sex steroid synthesis includes administering a therapeutically effective amount of an inhibitor of 5 α -reductase.

3. The method of Claim 1 wherein said inhibition of sex steroid synthesis includes administering a therapeutically effective amount of an inhibitor of 17 β -hydroxysteroid dehydrogenase.

4. The method of Claim 1 wherein said inhibition of sex steroid synthesis includes administering a therapeutically effective amount of an inhibitor of 5 α -reductase and an inhibitor of 17 β -hydroxysteroid dehydrogenase.

5. The method according to Claim 1 further comprising the step of administering a therapeutically effective amount of an antiestrogen.
6. The method according to Claim 2 further comprising the step of administering a therapeutically effective amount of an antiestrogen.
7. The method according to Claim 3 further comprising the step of administering a therapeutically effective amount of an antiestrogen.
8. The method according to Claim 4 further comprising the step of administering a therapeutically effective amount of an antiestrogen.
9. The method of Claim 1 wherein said inhibition includes the step of administering a mixture of sex steroid inhibitors which together are capable of inhibition the function of 5α -reductase and the function of 17β -hydroxysteroid dehydrogenase and the function of 3β -hydroxysteroid dehydrogenase.
10. The method according to Claim 9 further comprising the step of administering a therapeutically effective amount of an antiestrogen.
11. The method of Claim 1 further comprising the step of inhibiting testicular hormonal secretion of said warm-blooded animal.
12. The method of Claim 11 wherein said testicular hormonal secretion is inhibited by at least one technique selected from the group consisting of surgical castration, administration of an antagonist of luteinizing hormone releasing hormone, and administration of an agonist of luteinizing hormone releasing hormone.


13. The method of Claim 1 wherein said inhibition of sex steroid synthesis includes the step of administering an inhibitor of 5 α -reductase activity, and wherein said method further includes inhibiting the testicular hormonal secretion of said warm-blooded animal.
14. The method of Claim 2 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
15. The method of Claim 3 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
16. The method of Claim 4 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
17. The method of Claim 5 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
18. The method of Claim 6 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
19. The method of Claim 7 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
20. The method of Claim 8 which includes inhibiting testicular hormone secretion of said warm-blooded animal.

21. The method of Claim 9 which includes inhibiting testicular hormone secretion of said warm-blooded animal.


22. The method of Claim 10 which includes inhibiting testicular hormone secretion of said warm-blooded animal.

23. The method of Claim 1 wherein said inhibitor of sex steroid formation is selected from the group consisting of:


N-butyl, N-methyl-11-(16' α -chloro-3',17' β -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 139").




N-n-butyl-N-methyl-11-(16' α -chloro-3',17' α -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 170")




N-n-butyl-N-methyl-11-(16' α -bromo-3',17' α -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 171")



24. The method of Claim 1 wherein said antiandrogen is represented by the formula:




wherein the dotted line represents an optional double bond; x is 0, R¹ is [-CH₂-]_y (with y being an integer from 4 to 20), R^{17(α)} is hydrogen, lower alkyl or a moiety which together with R^{17(β)} forms:




III

wherein R^{17(β)} is selected from the group consisting of hydrogen, hydroxyl, lower alkyl, acyloxy, and a moiety which, together with R^{17(α)} forms:




III

25. The method of Claim 1 wherein said antiandrogen is represented by the general formula:




wherein the dotted lines represent optional double bonds; wherein R^{10} is hydrogen or lower alkyl, R^{13} is absent, hydrogen or methyl in β position,

$R^{17(\alpha)}$ is selected from the group consisting of hydrogen, hydroxyl, lower alkanoyloxy, lower alkyl, lower alkenyl, lower alkynyl, halo(lower)alkyl, halo(lower)alkenyl, halo(lower)alkynyl and fluoro-substituted aromatic ring, and a moiety which, together with $R^{17(\beta)}$ forms




III

$R^{17(\beta)}$ is selected from the group consisting of hydroxyl, (C_1-C_{10}) alkanoyloxy, (C_2-C_9) alkenoyloxy, (C_2-C_9) alkynoyloxy, aroyloxy, alkanoyloxy, cycloalkenoyloxy, 1-alkyloxy-alkyloxy, 1-alkyloxycycloalkyloxy, alkyl-




silyloxy, carboxyl, alkanoyl and a moiety which together with R¹⁷ forms



III

26. The method of Claim 1 wherein said antiandrogen is represented by the general formula:



VII

wherein the AB-ring junction is trans, the dotted lines represent optional pi bonds; wherein y is an integer from 4 to 20, wherein L is selected from the group consisting of -CONR^a-, -CSNR^a-, -NR^bCO-, -NR^bCS- and -CH²- (R^a and R^b being hydrogen or methyl) and G is selected from the group consisting of n-propyl, n-butyl, n-pentyl and haloalkyl.

27. A method for treatment of prostate cancer in a human or other warm-blooded animal in need of such treatment. Said method comprising the steps of inhibiting the sex steroid formation by administering a therapeutically effect of amount of an inhibitor of sex steroid formation

having, as part of the molecular structure, a sex-steroid nucleus and, as another part of its molecular structure, at least one side chain represented by the formula:

$-R^1[-B-R^2-]_x L-G$ wherein:

x is an integer from 0 to 6, wherein at least one of L and G is a polar moiety distanced from said ring carbon by at least three intervening atoms, and wherein:

R^1 and R^2 are independently either absent or selected from the group consisting of straight- or branched-chain alkylene, straight- or branched chain alkynylene, straight- or branched-chain alkenylene, phenylene and fluoro-substituted analogs of the foregoing;

B is either absent or selected from the group consisting of $-O-$, $-S-$, $-Se-$, $-SO-$, $-SO_2-$, $-NR^3-$, $-SiR^3_2-$, $-CR^3OR^3-$, $-NR^3CO-$, $-NR^3CS-$, $-CONR^3-$, $-CSNR^3-$, $-COO-$, $-COS-$, $-SCO-$, $-CSS-$, $-SCS-$, $-OCO-$ and phenylene (R^3 being hydrogen or lower alkyl);

L is either a moiety which together with G, forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of lower alkyl, $-CONR^4-$, $-CSNR^4-$, $-NR^4CO-$, $-NR^4CS-$, $-NR^4CONR^4-$

NR^6

$-NR^4C-NR^4-$, $-SO_2NR^4-$, $-CSS-$, $-SCS-$, $-(NO)R^4-$, $-(PO)R^4-$, $-NR^4COO-$, $-NR^4SO_2-$, $-O-$, $-NR^4-$, $-S-$, $-SO-$ and $-SO_2-$ (R^4 and R^5 being independently selected from the group consisting of hydrogen and lower alkyl; and R^6 being selected from the group consisting of hydrogen, nitrile and nitro); and

G is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, (C_3-C_{10}) cycloalkyl, bromo(lower)alkyl, chloro(lower)alkyl, fluoro(lower)alkyl, cyano(lower)alkyl, carboxy(lower)alkyl, (lower)alkoxycarbonyl(lower)alkyl, (C_6-C_{10}) aryl, $(C_7-C_{11})-$

arylalkyl, di(lower)alkylamino(lower)alkyl, fluoro-substituted analogs of the foregoing and a moiety which together with L forms a heterocyclic ring having at least one nitrogen atom; said method of treating prostate cancer further comprising the step of administering a therapeutically effective amount of an antiandrogen)

28. The method according to Claim 27 wherein the same method includes the step of administering a therapeutically effective amount of an inhibitor of 5 α -reductase

29. The method of Claim 27 wherein said inhibition of sex steroid synthesis includes administering a therapeutically effective amount of an inhibitor of 17 β -hydroxysteroid dehydrogenase.

30. The method of Claim 27 wherein said inhibition of sex steroid synthesis includes administering a therapeutically effective amount of an inhibitor of 5 α -reductase and an inhibitor of 17 β -hydroxysteroid dehydrogenase.

31. The method according to Claim 27 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

32. The method according to Claim 28 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

33. The method according to Claim 29 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

34. The method according to Claim 30 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

35. The method of Claim 27 wherein said inhibition includes the step of administering a mixture of sex steroid inhibitors which together are capable of inhibition the function of 5α -reductase and the function of 17β -hydroxysteroid dehydrogenase and the function of 3β -hydroxysteroid dehydrogenase.

36. The method according to Claim 35 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

37. The method of Claim 27 further comprising the step of inhibiting testicular hormonal secretion of said warm-blooded animal.

38. The method of Claim 37 wherein said testicular hormonal secretion is inhibited by at least one technique selected from the group consisting of surgical castration, administration of an antagonist of luteinizing hormone releasing hormone, and administration of an agonist of luteinizing hormone releasing hormone.

39. The method of Claim 27 wherein said inhibition of sex steroid synthesis includes the step of administering an inhibitor of 5α -reductase activity, and wherein said method further includes inhibiting the testicular hormonal secretion of said warm-blooded animal.

40. The method of Claim 28 which includes inhibiting the testicular hormonal secretion of said warm-blooded animal.

41. The method of Claim 29 which includes inhibiting the testicular hormonal secretion of said warm-blooded animal.

42. The method of Claim 30 which includes inhibiting the testicular hormonal secretion of said warm-blooded animal.

43. The method of Claim 31 which includes inhibiting the testicular hormonal secretion of said warm-blooded animal.

44. The method of Claim 32 which includes inhibiting the testicular hormonal secretion of said warm-blooded animal.

45. The method of Claim 33 which includes inhibiting the testicular hormonal secretion of said warm-blooded animal.

46. The method of Claim 34 which includes inhibiting the testicular hormonal secretion of said warm-blooded animal.


47. The method of Claim 35 which includes inhibiting the testicular hormonal secretion of said warm-blooded animal.

48. The method of Claim 36 which includes inhibiting the testicular hormonal secretion of said warm-blooded animal.


49. The method of Claim 27 wherein said inhibitor of sex steroid formation is selected from the group consisting of:

SUBSTITUTE SHEET


N-butyl, N-methyl-11-(16' α -chloro-3',17' β -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 139").




N-n-butyl-N-methyl-11-(16' α -chloro-3',17' α -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 170")




N-n-butyl-N-methyl-11-(16' α -bromo-3',17' α -dihydroxy-estra-1',3',5'(10')-trien-7' α -yl) undecanamide ("EM 171")



50. The method of Claim 27 wherein said antiandrogen is represented by the formula:




wherein the dotted line represents an optional double bond; x is 0, R¹ is $[-CH_2-]_y$ (with y being an integer from 4 to 20), R¹⁷(α) is hydrogen, lower alkyl or a moiety which together with R¹⁷(β) forms:




III

wherein R¹⁷(β) is selected from the group consisting of hydrogen, hydroxyl, lower alkyl, acyloxy, and a moiety which, together with R¹⁷(α) forms:




III

51. The method of Claim 27 wherein said antiandrogen is represented by the general formula:



wherein the dotted lines represent optional double bonds; wherein R¹⁰ is hydrogen or lower alkyl, R¹³ is absent, hydrogen or methyl in β position,


R^{17(α)} is selected from the group consisting of hydrogen, hydroxyl, lower alkanoyloxy, lower alkyl, lower alkenyl, lower alkynyl, halo(lower)alkyl, halo(lower)alkenyl, halo(lower)alkynyl and fluoro-substituted aromatic ring, and a moiety which, together with R^{17(β)} forms



III


R^{17(β)} is selected from the group consisting of hydroxyl, (C₁-C₈) alkanoyloxy, (C₂-C₈) alkenoyloxy, (C₂-C₈) alkynoyloxy, aryloxy, alkanoyloxy, cycloalkanyloxy, 1-alkyloxy-alkyloxy, 1-alkyloxycycloalkyloxy, alkyl-

silyloxy, carboxyl, alkanoyl and a moiety which together with R¹⁷ forms



III


52. The method of Claim 27 wherein said antiandrogen is represented by the general formula:



wherein the AB-ring junction is trans, the dotted lines represent optional pi bonds; wherein y is an integer from 4 to 20, wherein L is selected from the group consisting of $-\text{CONR}^a-$, $-\text{CSNR}^a-$, $-\text{NR}^b\text{CO}-$, $-\text{NR}^b\text{CS}-$ and $-\text{CH}_2-$ (R^a and R^b being hydrogen or methyl) and G is selected from the group consisting of n-propyl, n-butyl, n-pentyl and haloalkyl.

53. The method of Claim 27 wherein the molecular structure of said antiandrogen is an androgen nucleus substitute data ring carbon with at least one side chain represented by the formula $-\text{R}^1[-\text{B}-\text{R}^2-] ? \text{L-G}$.

54. The method of Claim 27 wherein said inhibitor of sex steroid formation has, as part of its molecular structure, a substituted or unsubstituted estrogen nucleus general structure I



wherein the dotted lines represent optional pi bonds and wherein said compound includes as another part of its molecular structure at least one side chain

substituent onto a ring carbon of said general structure I represented by the formula $-R^1[-B-R^2-]_x L-G$ wherein at least one of said side chains is substituted at a ring carbon selected from the group consisting of carbon 14, carbon 15, carbon 16 and carbon 17, wherein:

x is an integer from 0 to 6, wherein at least one of L and G is a polar moiety distanced from said ring carbon by at least three intervening atoms, and wherein:

R^1 and R^2 are independently either absent or selected from the group consisting of straight- or branched-chain alkylene, straight- or branched-chain alkynylene, straight- or branched-chain alkenylene, phenylene, and fluoro-substituted analogs of the foregoing;

B is either absent or selected from the group consisting of -O-, -S-, -Se-, -SO-, -SO₂-, -NR³-, -SiR³-, -CR³OR³-, -NR³CO-, -NR³CS-, -CONR³-, -CSNR³-, -COO-, -COS-, -SCO-, -CSS-, -SCS-, -OCO- and phenylene (R^3 being hydrogen or lower alkyl);


L is either a moiety which together with G, forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of lower alkyl, -CONR⁴-, -CSNR⁴-, -NR⁴CO-, -NR⁴CS-, -NR⁴CONR⁴-,
NR⁴

-NR⁴C-NR⁴-, -SO₂NR⁴-, -CSS-, -SCS-, -(NO)R⁴-, -(PO)R⁴-, -NR⁴COO-, -NR⁴SO₂-, -O-, -NR⁴-, -S-, -SO- and -SO₂- (R^4 and R^5 being independently selected from the group consisting of hydrogen and lower alkyl; and R^6 being selected from the group consisting of hydrogen, nitrile and nitro); and

G is either a moiety which together with L forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, (C₃-C₇)cyclo-


alkyl, bromo(lower)alkyl, chloro(lower)alkyl, fluoro(lower)alkyl, iodo(lower)alkyl, cyano(lower)alkyl, carboxy(lower)alkyl, (lower)alkoxycarbonyl(lower)alkyl, (C₆-C₁₀)aryl, (C₇-C₁₁)arylalkyl, di(lower)alkylamino(lower)alkyl, fluoro-substituted analogs of the foregoing.

55. The method of Claim 27 wherein said inhibitor of sex steroid formation has, as part of its molecular structure, a substituted or unsubstituted estrogen nucleus of general structure I.




wherein the dotted lines represent optional pi bonds; and wherein said compound includes, as another part of its molecular structure, at least one substitution of a side chain represented by the formula $-R^1[-B-R^1] L-G$ onto a ring carbon of said general structure I in at least one position selected from the group consisting of 7, 14, 15, 16 and 17, wherein said general structure I further includes at least one substitution selected from the group consisting of 15-halo, 16-halo, a 15,16 bridge atom (preferably carbon), a 14,15 bridge atom (preferably oxygen), 16-pi-bonded lower alkyl.

56. The method of Claim 27 wherein said antiandrogen is represented by the formula:




wherein the dotted line represents an optional double bond; x is 0, R^1 is $[-CH_2-]_y$ (with y being an integer from 4 to 20), $R^{17(\alpha)}$ is hydrogen, lower alkyl or a moiety which together with $R^{17(\beta)}$ forms:



III


wherein $R^{17(\beta)}$ is selected from the group consisting of hydrogen, hydroxyl, lower alkyl, acyloxy, and a moiety which, together with $R^{17(\alpha)}$ forms:




III

SUBSTITUTE SHEET

57. The method of Claim 27 wherein said antiandrogen is represented by the general formula:




wherein the dotted lines represent optional double bonds; wherein R^{10} is hydrogen or lower alkyl, R^{13} is absent, hydrogen or methyl in β position, $R^{17(\alpha)}$ is selected from the group consisting of hydrogen, hydroxyl, lower alkanoyloxy, lower alkyl, lower alkenyl, lower alkynyl, halo(lower)alkyl, halo(lower)alkenyl, halo(lower)alkynyl and fluoro-substituted aromatic ring, and a moiety which, together with $R^{17(\beta)}$ forms



III


$R^{17(\beta)}$ is selected from the group consisting of hydroxyl, (C_1-C_{20}) alkanoyloxy, (C_2-C_{20}) alkenoyloxy, (C_2-C_{20}) alkynoyloxy, arcyloxy, alkanoyloxy, cycloalkenoyloxy, 1-alkyloxy-alkyloxy, 1-alkyloxycycloalkyloxy, alkyl-

silyloxy, carboxyl, alkanoyl and a moiety which together with R¹⁷ forms



III

58. The method of Claim 27 wherein said antiandrogen is represented by the general formula:



wherein the AB-ring junction is trans, the dotted lines represent all pi bonds; wherein y is an integer from 4 to 20, wherein L is selected from the group consisting of -CONR^a-, -CSNR^a-, -NR^bCO-, -NR^bCS- and (R^a and R^b being hydrogen or methyl) and G is selected from the group consisting of n-propyl, n-butyl, n-pentyl and haloalkyl.

59. A method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of administering a therapeutically effective amount of an inhibitor of 5 α -reductase activity and administering a therapeutically effective amount of an antiandrogen.

60 The method of Claim 59 further comprising administering a therapeutically effective amount of at least one compound selected from the group consisting of an antiestrogen, an inhibitor of aromatase activity, an inhibitor of 17 β -hydroxysteroid dehydrogenase activity, and inhibitor of testicular hormonal secretion.

61. The method of Claim 59 where testicular hormone secretion is blocked by surgical castration.

62. The method of Claim 55 wherein said side chain is substituted onto said estrogen nucleus in a position selected from the group consisting of 7 α , 15 α and 17 α .

63. A method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of administering a therapeutically effective amount of an inhibitor of 17 β -HSD and administering a therapeutically effective amount of an antiandrogen.

64. A method of Claim 63 further comprising administering a therapeutically effective amount of at least one compound selected from the group consisting of an antiestrogen, an inhibitor of 5 α -reductase activity, an LHRH agonist and an LHRH antagonist.

65. The method of Claim 63 where testicular hormone secretion is blocked by surgical castration.

66. The method of Claim 64 where testicular hormone secretion is blocked by surgical castration.

67. A method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of administering an effective amount of an antiandrogen and at least one compound selected from the groups consisting of inhibitors of extragonadal and extraadrenal sex steroid formation.

68. The method of Claim 67 where testicular steroid secretion is blocked by surgical castration.

69. The method of Claim 67 where serum said inhibition of sex steroid synthesis includes administering a therapeutically effective amount of an inhibitor of 5 α -reductase.

70. The method of Claim 67 where serum said inhibition of sex steroid synthesis includes administering a therapeutically effective amount of an inhibitor of 17 β -hydroxysteroid dehydrogenase.

71. The method of Claim 67 wherein said inhibition of sex steroid synthesis includes administering a therapeutically effective amount of an inhibitor of 5 α -reductase and an inhibitor of 17 β -hydroxysteroid dehydrogenase.

72. The method according to Claim 67 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

73. The method according to Claim 69 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

74. The method according to Claim 70 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

75. The method according to Claim 71 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

76. The method of Claim 67 further comprising the step of inhibiting testicular hormonal secretion of said warm-blooded animal.

77. The method of Claim 76 wherein said testicular hormonal secretion is inhibited by at least one technique selected from the group consisting of surgical castration, administration of an antagonist of luteinizing hormone releasing hormone, and administration of an agonist of luteinizing hormone releasing hormone.

78. The method of Claim 67 wherein said inhibition of sex steroid synthesis includes the step of administering an inhibitor of 5 α -reductase activity, and wherein said method further includes inhibiting the testicular hormonal secretion of said warm-blooded animal.


79. The method of Claim 69 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
80. The method of Claim 70 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
81. The method of Claim 71 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
82. The method of Claim 72 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
83. The method of Claim 73 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
84. The method of Claim 74 which includes inhibiting testicular hormone secretion of said warm-blooded animal.
85. The method of Claim 75 which includes inhibiting testicular hormone secretion of said warm-blooded animal.

86. A kit for treating prostate cancer, said kit including a plurality of separate containers, each containing at least one active ingredient useful in a combination therapy for treatment of prostate cancer (with or without additional pharmaceutical carrier or diluent) and each container containing at least one active ingredient which is not contained in any other container in said kit, wherein one of said containers contains an antiandrogen and another of said containers contains an active ingredient selected from the group consisting of an inhibitor of 5 α -reductase activity and an inhibitor capable of blocking formation of natural sex steroids from dehydroepiandrosterone and from 4-androstenedione in peripheral tissues.

87. The kit of Claim 86 wherein said kit further comprises at least one separate container containing an active ingredient selected from the group consisting of an inhibitor of testicular hormonal secretion, an inhibitor of aromatase, an inhibitor of 17 β -hydroxysteroid dehydrogenase and an inhibitor of 3 β -hydroxysteroid dehydrogenase.

88. A kit for treating prostate cancer, said kit including a plurality of separate containers, each containing at least one active ingredient useful in a combination therapy for treatment of prostate cancer (with or without additional pharmaceutical carrier or diluent) and each container containing at least one active ingredient which is not contained in any other container in said kit, wherein one of said containers contains an inhibitor of sex steroid formation and another of said containers contains (with or without carrier or diluent) an antiandrogen

having, as part of its molecular structure, a substituted or unsubstituted androgenic nucleus of the formula:



and having as another part of its molecular structure, at least one side chain represented by the formula:

$-R^1[-B-R^2-]_x L-G$ said chain being substituted onto said androgenic nucleus at a position selected from the group consisting of 6, 7, 14, 15, 16 and 17, wherein:

x is an integer from 0 to 6, wherein at least one of L and G is a polar moiety distanced from said ring carbon by at least three intervening atoms, and wherein:

R^1 and R^2 are independently either absent or selected from the group consisting of straight- or branched-chain alkylene, straight- or branched chain alkynylene, straight- or branched-chain alkenylene, phenylene and fluoro-substituted analogs of the foregoing;

B is either absent or selected from the group consisting of $-O-$, $-S-$, $-Se-$, $-SO-$, $-SO_2-$, $-NR^3-$, $-SiR^3-$, $-CR^3OR^3-$, $-NR^3CO-$, $-NR^3CS-$, $-CONR^3-$, $-CSNR^3-$, $-COO-$, $-COS-$, $-SCO-$, $-CSS-$, $-SCS-$, $-OCO-$ and phenylene (R^3 being hydrogen or lower alkyl);

SUBSTITUTE SHEET

L is either a moiety which together with G, forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of lower alkyl, -CONR^a-, -CSNR^a-, -NR^sCO-, -NR^sCS-, -NR^sCONR^a-

NR^s
||
-NR^sC-NR^a-, -SO₂NR^a-, -CSS-, -SCS-, -(NO)R^a-, -(PO)R^a-, -NR^sCOO-,
-NR^sSO₂-, -O-, -NR^a-, -S-, -SO- and -SO₂- (R^a and R^s being independently selected from the group consisting of hydrogen and lower alkyl; and R^s being selected from the group consisting of hydrogen, nitrile and nitro);
and


G is either a moiety which together with L forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, (C₃-C₆)cycloalkyl, bromo(lower)alkyl, chloro(lower)alkyl, fluoro(lower)alkyl, cyano(lower)alkyl, carboxy(lower)alkyl, (lower)alkoxycarbonyl(lower)alkyl, (C₆-C₁₀)aryl, (C₇-C₁₁)arylalkyl, di(lower)alkylamino(lower)alkyl, fluoro-substituted analogs of the foregoing and a moiety which together with L forms a heterocyclic ring having at least one nitrogen atom;

89. The kit of Claim 88 wherein said kit further comprises at least one separate container containing an active ingredient selected from the group consisting of an inhibitor of testicular hormonal secretion, an inhibitor of aromatase, an inhibitor of 17β-hydroxysteroid dehydrogenase and an inhibitor of 3β-hydroxysteroid dehydrogenase.

90. A pharmaceutical composition for treatment of prostate cancer comprising an antiandrogen (with or without a pharmaceutical diluent or carrier) and at least one active ingredient selected from the group consisting of an inhibitor of 5 α -reductase activity and an inhibitor capable of blocking formation of natural sex steroids from dehydroepiandrosterone and from 4-androstenedione in peripheral tissues.

91. The pharmaceutical composition of Claim 90 further comprising at least one active ingredient selected from the group consisting of an inhibitor of testicular hormonal secretions, an inhibitor of aromatase, an inhibitor of 17 β -hydroxysteroid dehydrogenase and an inhibitor of 3 β -hydroxysteroid dehydrogenase.

92. A pharmaceutical composition for treatment of prostate cancer comprising an inhibitor of sex steroid formation and an antiandrogen having, as part of its molecular structure, a substituted or unsubstituted androgenic nucleus of the formula:



and having as another part of its molecular structure at least one side chain represented by the formula:

$-R^1[-B-R^2-]_x$ L-G said chain being substituted onto said androgenic nucleus at a position selected from the group consisting of 6, 7, 14, 15, 16 and 17, wherein:

x is an integer from 0 to 6, wherein at least one of L and G is a polar moiety distanced from said ring carbon by at least three intervening atoms, and wherein:

R^1 and R^2 are independently either absent or selected from the group consisting of straight- or branched-chain alkylene, straight- or branched chain alkynylene, straight- or branched-chain alkenylene, phenylene and fluoro-substituted analogs of the foregoing;

B is either absent or selected from the group consisting of -O-, -S-, -Se-, -SO-, -SO₂-, -NR³-, -SiR³-, -CR³OR³-, -NR³CO-, -NR³CS-, -CONR³-, -CSNR³-, -COO-, -COS-, -SCO-, -CSS-, -SCS-, -OCO- and phenylene (R^3 being hydrogen or lower alkyl);

L is either a moiety which together with G, forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of lower alkyl, -CONR⁴-, -CSNR⁴-, -NR⁴CO-, -NR⁴CS-, -NR⁴CONR⁴-

NR⁴

-NR⁴C-NR⁴-, -SO₂NR⁴-, -CSS-, -SCS-, -(NO)R⁴-, -(PO)R⁴-, -NR⁴COO-, -NR⁴SO₂-, -O-, -NR⁴-, -S-, -SO- and -SO₂- (R^4 and R^5 being independently selected from the group consisting of hydrogen and lower alkyl; and R^6 being selected from the group consisting of hydrogen, nitrile and nitro); and

SUBSTITUTE SHEET

G is either a moiety which together with L forms a heterocyclic ring having at least one nitrogen atom or is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, (C₁-C₇)cycloalkyl, bromo(lower)alkyl, chloro(lower)alkyl, fluoro(lower)alkyl, cyano(lower)alkyl, carboxy(lower)alkyl, (lower)alkoxycarbonyl(lower)alkyl, (C₆-C₁₀)aryl, (C₁-C₁₁)arylalkyl, di(lower)alkylamino(lower)alkyl, fluoro-substituted analogs of the foregoing and a moiety which together with L forms a heterocyclic ring having at least one nitrogen atom.

93. The pharmaceutical composition of Claim 92 wherein said composition further comprises at least one active ingredient selected from the group consisting of an inhibitor of testicular hormonal secretions, an inhibitor of aromatase activity an inhibitor of 17 β -hydroxysteroid dehydrogenase and an inhibitor of 3 β -hydroxysteroid dehydrogenase.

94. The method of claim 27 wherein said antianδrogen is selected from the group consisting of Flutamide, Anandron, Casodex and Androcur.

95. A method for treatment of prostate cancer in a human or other warm-blooded animal in need of such treatment. Said method comprising the steps of inhibiting the sex steroid formation by administering a therapeutically effect of amount of an inhibitor of sex steroid formation having, as part of the molecular structure, a sex-steroid nucleus and, as another part of its molecular structure, at least one side chain represented by the formula:

$-R^1[-B-R^2-]_x L-G$ wherein:

x is an integer from 0 to 6, wherein at least one of L and G is a polar moiety distanced from said ring carbon by at least three intervening atoms, and wherein:

R^1 and R^2 are independently either absent or selected from the group consisting of straight- or branched-chain alkylene, straight- or branched chain alkynylene, straight- or branched-chain alkenylene, phenylene and fluoro-substituted analogs of the foregoing;

B is either absent or selected from the group consisting of -O-, -S-, -Se-, -SO-, -SO₂-, -NR³-, -SiR³-, -CR³OR³-, -NR³CO-, -NR³CS-, -CONR³-, -CSNR³-, -COO-, -COS-, -SCO-, -CSS-, -SCS-, -OCO- and phenylene (R^3 being hydrogen or lower alkyl);

L is either a moiety which together with G, forms a heterocyclic ring having at least one nitrogen atom or is selected from the group

consisting of lower alkyl, $-\text{CONR}^A-$, $-\text{CSNR}^A-$, $-\text{NR}^B\text{CO}-$, $-\text{NR}^B\text{CS}-$, $-\text{NR}^A\text{CONR}^B-$,
 NR^A

$-\text{NR}^B\text{C}-\text{NR}^A-$, $-\text{SO}_2\text{NR}^A-$, $-\text{CSS}-$, $-\text{SCS}-$, $-(\text{NO})\text{R}^A-$, $-(\text{PO})\text{R}^A-$, $-\text{NR}^B\text{COO}-$,
 $-\text{NR}^B\text{SO}_2-$, $-\text{O}-$, $-\text{NR}^A-$, $-\text{S}-$, $-\text{SO}-$ and $-\text{SO}_2-$ (R^A and R^B being independently
 selected from the group consisting of hydrogen and lower alkyl; and R^B
 being selected from the group consisting of hydrogen, nitrile and nitro);
 and

G is selected from the group consisting of hydrogen, lower alkyl,
 lower alkenyl, lower alkynyl, $(\text{C}_3-\text{C}_7)\text{cycloalkyl}$, bromo(lower)alkyl,
 chloro(lower)alkyl, fluoro(lower)alkyl, cyano(lower)alkyl, carboxy-
 (lower)alkyl, (lower)alkoxycarbonyl(lower)alkyl, $(\text{C}_6-\text{C}_{10})\text{aryl}$, $(\text{C}_7-\text{C}_{11})$ -
 arylalkyl, di(lower)alkylamino(lower)alkyl, fluoro-substituted analogs of
 the foregoing and a moiety which together with L forms a heterocyclic
 ring having at least one nitrogen atom; said method of treating prostate
 cancer further comprising the step of administering a therapeutically
 effective amount of an inhibitor of 5α -reductase.

96. A method for treating prostate cancer
 in a human or other warm-blooded animal in need of such
 treatment, said method comprising the steps of
 inhibiting testicular hormonal secretion of said warm-
 blooded animal, and administering a pharmaceutically
 effective amount of a 5α reductase inhibitor.

97. The method of claim 96 wherein said method further comprises the step of administering a pharmaceutically effective amount of at least one compound selected from the group consisting of an antiestrogen, an inhibitor of 17 β -hydroxysteroid dehydrogenase and an inhibitor of aromatase activity.

98. A method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of inhibiting testicular hormonal secretion of said warm-blooded animal, and administering a pharmaceutically effective amount of an antiestrogen.

99. The method of claim 98 wherein said method further comprises the step of administering a pharmaceutically effective amount of an antiandrogen.

100. The method of claim 98 wherein said method further comprises the step of administering an effective amount of a 5 α -reductase inhibitor.

101. The method of claim 100 wherein said 5 α -reductase inhibitor is Proscar.


102. The method of claim 98 wherein said testicular hormonal secretion is inhibited by at least one technique selected from the group consisting of surgical castration, administration of an antagonist of luteinizing hormone releasing hormone, and administration of an agonist of luteinizing hormone releasing hormone.

103. A method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of administering a pharmaceutically effective amount of an antiestrogen in combination with a pharmaceutically effective amount of at least one compound selected from the group consisting of an antiandrogen, a 5 α -reductase inhibitor and a sex steroid inhibitor.

104. The method of claim 2, wherein said inhibitor of 5 α -reductase is Proscar.


105. The method of claim 28, wherein said inhibitor of 5 α -reductase is Proscar.

106. The method of claim 1 wherein said inhibitor of sex steroid formation is:



wherein R is either hydrogen or ethynyl.

107. The method of claim 27 wherein said inhibitor of sex steroid formation is:



wherein R is either hydrogen or ethynyl.

AMENDED CLAIMS

[received by the International Bureau on 26 December 1990 (26.12.90);
original claims 28,60,97,106 and 107 amended;
new claims 108-116 added; other claims unchanged
(6 pages)]

arylalkyl, di(lower)alkylamino(lower)alkyl, fluoro-substituted analogs of the foregoing and a moiety which together with I. forms a heterocyclic ring having at least one nitrogen atom; said method of treating prostate cancer further comprising the step of administering a therapeutically effective amount of an antiandrogen)

28. The method according to Claim 27 wherein the same method includes the step of administering a therapeutically effective amount of an inhibitor of 5 α -reductase

29. The method of Claim 27 wherein said inhibition of sex steroid synthesis includes administering a therapeutically effective amount of an inhibitor of 17 β -hydroxysteroid dehydrogenase.

30. The method of Claim 27 wherein said inhibition of sex steroid synthesis includes administering a therapeutically effective amount of an inhibitor of 5 α -reductase and an inhibitor of 17 β -hydroxysteroid dehydrogenase.

31. The method according to Claim 27 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

32. The method according to Claim 28 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

33. The method according to Claim 29 further comprising the step of administering a therapeutically effective amount of an antiestrogen.

59. A method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of administering a therapeutically effective amount of an inhibitor of 5 α -reductase activity and administering a therapeutically effective amount of an antiandrogen.

60. The method of Claim 59 further comprising administering a therapeutically effective amount of at least one compound selected from the group consisting of an antiestrogen, an inhibitor of aromatase activity, an inhibitor of 17 β -hydroxysteroid dehydrogenase activity, and inhibitor of testicular hormonal secretion.

61. The method of Claim 59 where testicular hormone secretion is blocked by surgical castration.

62. The method of Claim 55 wherein said side chain is substituted onto said estrogen nucleus in a position selected from the group consisting of 7 α , 15 α and 17 α .

63. A method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of administering a therapeutically effective amount of an inhibitor of 17 β -HSD and administering a therapeutically effective amount of an antiandrogen..

97. The method of claim 96 wherein said method further comprises the step of administering a pharmaceutically effective amount of at least one compound selected from the group consisting of an antiestrogen, an inhibitor of 17 β -hydroxysteroid dehydrogenase and an inhibitor of aromatase activity.


98. A method for treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of inhibiting testicular hormonal secretion of said warm-blooded animal, and administering a pharmaceutically effective amount of an antiestrogen.

99. The method of claim 98 wherein said method further comprises the step of administering a pharmaceutically effective amount of an antiandrogen.

100. The method of claim 98 wherein said method further comprises the step of administering an effective amount of a 5 α -reductase inhibitor.


101. The method of claim 100 wherein said 5 α -reductase inhibitor is Proscar.

106. The method of claim 1 wherein said inhibitor of sex steroid formation is:



wherein R is either hydrogen or ethynyl.

107. The method of claim 27 wherein said inhibitor of sex steroid formation is:



wherein R is either hydrogen or ethynyl.

108. The method of claim 95 wherein said method of treating prostate cancer further comprises the step of administering a therapeutically effective amount of a pharmaceutical selected from the group consisting of an
5 inhibitor of 17 β -hydroxysteroid dehydrogenase, an inhibitor of 3 β -hydroxysteroid dehydrogenase, and an inhibitor of aromatase.

109. The method of claim 100, wherein said method further comprises the step of administering a
10 pharmaceutically effective amount of an antiandrogen.

110. The method of claim 103, wherein said method further comprises the step of inhibiting testicular hormone secretion.

111. A method for treating or preventing prostate
15 cancer in a human or other warm-blooded animal in need of such treatment, said method comprising the step of administering to said warm-blooded animal a pharmaceutically effective amount of an antiestrogen.

112. The method of claim 59, wherein said
20 antiandrogen is selected from the group consisting of Flutamide, Anandron, Casodex and Androcur, and wherein the 5 α -reductase inhibitor is Prosoar.


113. The method of claim 1, wherein said sex steroid formation inhibitor is an aromatase inhibitor.

25 114. The method of claim 27, wherein said sex steroid formation inhibitor is an aromatase inhibitor.

115. The method of claim 67, wherein testicular hormonal secretion is blocked by treatment with an LHRH agonist or LHRH antagonist.




116. A method of treating prostate cancer in a human or other warm-blooded animal in need of such treatment, said method including the steps of administering a pharmaceutically effective amount of a 5 α -reductase inhibitor and at least one compound selected from the group consisting of an antiestrogen, an inhibitor of 17 β -hydroxy steroid dehydrogenase, an aromatase inhibitor, and an inhibitor of testicular hormonal secretion.

1/1



INTERNATIONAL SEARCH REPORT

International Application No PCT/CA 90/00212

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) * According to International Patent Classification (IPC) or to both National Classification and IPC IPC ⁵ : A 61 K 31/565, A 61 K 31/575, A 61 K 31/57, // (A 61 K 31/575, 31:57, 31:565), (A 61 K 31/57, 31:565)																				
II. FIELDS SEARCHED <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Minimum Documentation Searched ?</div> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%; border-bottom: 1px solid black;">Classification System</td> <td style="border-bottom: 1px solid black;">Classification Symbols</td> </tr> <tr> <td style="padding: 5px;">IPC⁵</td> <td style="padding: 5px;">A 61 K</td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched *</div>			Classification System	Classification Symbols	IPC ⁵	A 61 K														
Classification System	Classification Symbols																			
IPC ⁵	A 61 K																			
III. DOCUMENTS CONSIDERED TO BE RELEVANT * <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 10%;">Category *</th> <th style="width: 70%;">Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²</th> <th style="width: 20%;">Relevant to Claim No. ¹³</th> </tr> </thead> <tbody> <tr> <td style="text-align: center; vertical-align: top;">X</td> <td style="vertical-align: top;"> WO, A, 8601105 (LABRIE) 27 February 1986 see the whole document; especially page 12, line 12 - page 19, line 3; claims; abstract cited in the application -- </td> <td style="text-align: center; vertical-align: top;">86,87,89-91</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">X</td> <td style="vertical-align: top;"> LU, A, 83545 (SCHERING) 11 October 1981 see the whole document -- </td> <td style="text-align: center; vertical-align: top;">86,87,89-91</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">X</td> <td style="vertical-align: top;"> WO, A, 8705216 (SCHERING) 11 September 1987 see the whole document -- </td> <td style="text-align: center; vertical-align: top;">86,87,89-91</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">Y</td> <td style="vertical-align: top;"> EP, A, 0285383 (MERCK) 5 October 1988 see the whole document -- </td> <td style="text-align: center; vertical-align: top;">86-93</td> </tr> <tr> <td colspan="3" style="text-align: center; padding-top: 10px;">./.</td> </tr> </tbody> </table>			Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³	X	WO, A, 8601105 (LABRIE) 27 February 1986 see the whole document; especially page 12, line 12 - page 19, line 3; claims; abstract cited in the application --	86,87,89-91	X	LU, A, 83545 (SCHERING) 11 October 1981 see the whole document --	86,87,89-91	X	WO, A, 8705216 (SCHERING) 11 September 1987 see the whole document --	86,87,89-91	Y	EP, A, 0285383 (MERCK) 5 October 1988 see the whole document --	86-93	./.		
Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³																		
X	WO, A, 8601105 (LABRIE) 27 February 1986 see the whole document; especially page 12, line 12 - page 19, line 3; claims; abstract cited in the application --	86,87,89-91																		
X	LU, A, 83545 (SCHERING) 11 October 1981 see the whole document --	86,87,89-91																		
X	WO, A, 8705216 (SCHERING) 11 September 1987 see the whole document --	86,87,89-91																		
Y	EP, A, 0285383 (MERCK) 5 October 1988 see the whole document --	86-93																		
./.																				
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 50%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div>																				
IV. CERTIFICATION <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;"> Date of the Actual Completion of the International Search <div style="text-align: center;">19th September 1990</div> </td> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;"> Date of Mailing of this International Search Report <div style="text-align: center;">26. 10. 90</div> </td> </tr> <tr> <td style="border-bottom: 1px solid black; padding: 5px;"> International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div> </td> <td style="border-bottom: 1px solid black; padding: 5px;"> Signature of Authorized Officer <div style="text-align: center;">  Nathalie Weinberg </div> </td> </tr> </table>			Date of the Actual Completion of the International Search <div style="text-align: center;">19th September 1990</div>	Date of Mailing of this International Search Report <div style="text-align: center;">26. 10. 90</div>	International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: center;">  Nathalie Weinberg </div>														
Date of the Actual Completion of the International Search <div style="text-align: center;">19th September 1990</div>	Date of Mailing of this International Search Report <div style="text-align: center;">26. 10. 90</div>																			
International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: center;">  Nathalie Weinberg </div>																			

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
Y	DE, A, 3339295 (SCHERING) 17 May 1984 see the whole document --	86-93
Y	US, A, 4235893 (BRODIE et al.) 25 November 1980 see the whole document cited in the application	86-93
Y	Endocrine Research Communications, vol. 4, no. 2, 1977, Marcel Dekker, Inc., N.A. Musto et al.: "A novel affinity column for isolation of androgen binding protein from rat epididymis", pages 147- 157 see abstract; figure 1 --	86-93
Y	Steroids, vol. 40, no. 6, December 1982, A.J. Solo et al.: "7alpha-alkyltesto- sterone derivatives: synthesis and activity as androgens and as acromatase inhibitors", pages 603-614 see the whole article cited in the application --	86-93
Y	The Prostate, vol. 5, 1984, Alan R. Liss, Inc., M.-A. De Larminat et al.: "Synthesis and evaluation of immobilized androgens for affinity chromatography in the purification of nuclear androgen receptor", pages 123-140 see the whole article cited in the application --	86-93
L	World Patent Information, vol. 11, no. 1, 1989, CEC/WIPO, (GB), K.E.H. Göhring et al.: "A giant step for mankind ?", pages 5-10 see the whole article --	86-93
A	DE, A, 2529969 (ROUSSEL-UCLAF) 22 January 1976 see the whole document --	86-93
A	US, A, 4472382 (LABRIE et al.) 18 September 1984 see the whole document -----	86-93

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

INCOMPLETELY

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND 'SEARCHABLE'

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers ... **... because they relate to subject matter not required to be searched by this Authority, namely:

** Claim numbers 1-85, 94-116

Please see Rule 39.1 (iv): methods for treatment of the human or animal body by surgery or therapy as well as diagnostic methods.

2. ☒ Claim numbers 86-93 because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

SEE ANNEX

3. ☐ Claim numbers because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

- 1) A compound cannot be sufficiently characterised by its pharmacological profile or its mode of action as it is done by expressions like:

"antiandrogen"

"Inhibitor of 5 alpha- reductase activity"

"inhibitor capable of blocking formation of natural sexsteroids from dehydroepiandrosterone and from 4-androstenedione in peripheral tissues"

"inhibitor of testicular hormonal secretions"

"inhibitor of aromatase"

"inhibitor of 17-beta-hydroxysteroid dehydrogenase"

"inhibitor of 3-beta-hydroxysteroid dehydrogenase"

- 2) Reasons of economy (see Guidelines for Examination in the European Patent Office, part B, chapter III, paragraph 3.6) made it necessary to restrict the search. This is especially applicable for the formula given in the claims 88 and 92. The search was performed on base of the given examples.
- 3) The formula given in claim 88 and 92 contains no androgenic nucleus but an estrogenic nucleus.

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

CA 9000212

SA 37984

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 15/10/90. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE-A- 2529969		LU-A- 72892	31-05-76
		NL-A- 7507994	06-01-76
		SE-A- 7507353	05-01-76
		US-A- 3975413	17-08-76
<hr/>			
US-A- 4472382	18-09-84	FR-A, B 2465486	27-03-81
		AU-B- 572938	19-05-88
		AU-A- 3726985	09-05-85
		AU-B- 542765	14-03-85
		AU-A- 6256580	09-04-81
		BE-A- 885308	19-03-81
		JP-A, B, C 56055315	15-05-81
		JP-A- 61118324	05-06-86
		US-A- 4851386	25-07-89
		US-A- 4743589	10-05-88
		US-A- 4728640	01-03-88
		US-A- 4745102	17-05-88
<hr/>			